



Russian olive (*Elaeagnus angustifolia*) as a herbal healer

Zeinab Amiri Tehranizadeh¹, Ali Baratian², Hossein Hosseinzadeh^{3*}

¹ Department of Medicinal Chemistry, School of Pharmacy, Mashhad University of Medical Sciences, Mashhad, Iran

² Department of Pharmaceutics, School of Pharmacy, Mashhad University of Medical Sciences, Mashhad, Iran

³ Department of Pharmacodynamics and Toxicology, Pharmaceutical Research Center, School of Pharmacy, Mashhad University of Medical Sciences, Mashhad, Iran

Article Info



Article Type:
Review

Article History:

Received: 23 Apr. 2016
 Revised: 31 July 2016
 Accepted: 21 Aug. 2016
 ePublished: 24 Sep. 2016

Keywords:

Elaeagnus angustifolia
 Russian olive
 Inflammation
 Traditional remedy
 Flavonoid


Abstract

Introduction: *Elaeagnus* spp. is one in the family of riparian trees growing near the rivers or water corridors. In this family, *Elaeagnus angustifolia* (Russian olive) is famous because of its medical applications.

Methods: A comprehensive review was performed to extract the related data from published literature.

Results: Traditionally, it has been used as an analgesic, antipyretic and diuretic herbal medicine.

Conclusion: This review aimed to provide the most important documentary information on the active components of *Elaeagnus* spp. and their relation to the pharmacological properties and compare them with reported medicinal effects.



Introduction

Elaeagnus spp. (Plantae>Rosales>Elaeagnaceae>Elaeagnus) is in the family of riparian trees growing near rivers or water corridors.¹ It is mostly found in central Asia² including Iran,³ Uzbekistan,² Syria⁴ and north-west of China⁵ and exotically in river banks of central Spain,⁶ Canada⁷ and west of United-states.¹ The family Elaeagnaceae, consists of three genera and 50 species. *Elaeagnus angustifolia* (Russian olive) (Fig. 1) and *E. pungens* are partially adapted to the center of Asia. *Shepherdia canadensis* belongs to the Unites-states and Canada. *Hippophae rhamnoides* is endemic in Europe.⁸ In this family; *E. angustifolia* is famous for its medical benefits. Traditionally, it has been used as an analgesic, antipyretic and diuretic herbal medicine. A large number of compounds have been derived from Russian olive and made this plant a source of flavonoids, alkaloids, minerals and vitamins. Several experimental studies have been done and some advances in drug formulation and herbal medicine have been achieved. This review tries to gather the most important documentary information

on its active components and their relation to the Russian olive pharmacological properties and compare them with reported medicinal effects. Another review on this plant has been published recently, but we tried to check more articles with conflicting reports for writing better and more comprehensive systematic review.⁹

Active ingredients

The aqueous and non-aqueous extracts of *E. angustifolia* are full of medically significant active ingredients. The extracts contain a variety of compounds such as flavonoids and alkaloids, simple sugars and complicated sterols (Fig. 2).

Flavonoids

Flavonoids (which are commonly referred to vitamin P) are a large set of polyphenolic compounds with a benzo-γ-pyrone structure and are found exclusively in plants. They can be categorized in different classes like: flavones, flavonols, flavanones, flavanonols, isoflavones and flavan-3-ols.¹⁰

*Corresponding author: Hossein Hosseinzadeh, Email: hosseinzadehh@mums.ac.ir



© 2016 The Author(s). This work is published by BioImpacts as an open access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by-nc/4.0/>). Non-commercial uses of the work are permitted, provided the original work is properly cited.




Fig. 1. A brief review of *E. angustifolia*.




Fig. 2. Major constituents of *E. angustifolia*.

In the pulp of Russian olive, 4 glycosylated flavonoids¹⁰ identified: Quercetin 3,4'-O- β -D-diglucoside, Isorhamnetin-3-O- β -D-galactopyranoside, Quercetin 3-O- β -D-Galactopyranoside- 4'-O- β -D-glucopyranoside and Isorhamnetin 3-O- β -D-Galactopyranoside-4'-O- β -D-glucopyranoside.¹¹

Rutin glycosylated kampferols such as kaempferol 7-p-coumaroyl-3-D-glucoside, kaempferol-3-D-glucopyranoside (astragalin) and kaempferol 3-O-D-glucosido-41-p-coumaroyl-7-O-D- acyl galactoside are other available flavonols of the fruits.¹²

Rutin eleagnoside and kampferol are also available in the flavonoid fraction of the fruit. In acetone extract from the bark of Russian olive, (flavan-3-ol¹⁰) obtained two catechins as well: (+)-catechin and (-)-epicatechin.¹³

Glycosylated flavonoids have difficulty in absorption; af-

ter hydrolysis by lactase phlorizin hydrolase (LPH), they should transfer from Na⁺-dependent glucose cotransporter in small intestine. Mostly, the glycosylated flavonoids pass the colon and exit. For these reasons, *E. angustifolia* flavonoids may also exit the colon and have no benefits. So, it should be taken into account whether the amount of absorption is enough for medical benefits or not.¹⁰

However, they have different roles in plants; for example, they play a major role in oxidative stress response and are also used as growth regulators. But, talking about the exact amount of the flavonoids in each plant is something difficult. Recent studies showed that geographical location may influence the quantity and quality of flavonoids.¹⁰

Fatty acids

Fatty acids of this family first recognized by Obodovski and Devyatnin. Total lipid content of *E. angustifolia* varies between 0.8% in pericarp to 26% in seeds per mass.¹²

Fruit methanol extract from Russian olive contains different kinds of fatty acids. In an experiment by Kusova *et al.* oleic acid and linoleic acid made up 92.8% of the petroleum extract of the fruit accompanied by a low concentration of fat-soluble vitamins.¹⁴

In another study which published in the journal of the chemistry of natural compounds on the Russian olive seeds, six epoxy acids and five hydroxy acids were collected: C18:0, C18:1 and C18:2 diepoxy acids and C18-hydroxy acids with double bonds at the 9-C atoms.¹⁵

Palmitic acid (16:1) is also present in trace amount (3%-10%)¹⁶ in both seeds and fruits,^{16,18} but, it is not a chemotaxonomic compound for this tree.¹⁶

In general, flesh of the fruit is low in lipids but seeds and pericarps of Russian olive are saturated with these compounds. Triacylglycerol with linoleic acid is the main lipid of the seeds. Saturated free fatty acids (octadecanoic acid 1.63% and hexadecanoic acid 3.91%)¹⁷ are eminent; however, the highest concentration among free fatty acids belongs to essential linoleic acid (49.12%).¹⁷ These amounts are somehow identical to the pericarp, except for the amount of triacylglycerol which is lower (18%-30%) in the pericarp.¹⁶

The amount of fatty acids and their derivatives are different from tree to tree based on cultivation soil, tree age and planting area.¹⁶ However, what is important is whether the amount of fatty acids is commercially efficient or not; far seeds with 26% fatty acid per total mass, it seems applicable but for the rest (pericarp, leaves and flowers), it needs cost-benefit experiments.

Sterols

The most significant sterol of Russian olive is β -sitosterol.^{16,18} It is found mainly in seeds, leaves and tree branches. Elaeagnoside with a sterol-like structure is derived from the name of this family and present in the fruits of Russian olive.^{12,19}

Carbohydrates

Usually, free sugars are the first descriptors of the fruits. In Russian olive, fructose and glucose are the predominant monosaccharides. The percentage of weight of fructose and glucose to the total dried weight of the fruit is 32%-

34% and 23%-24% respectively.²⁰ In the ripened fruits there is no significant amount of sucrose; it is said that this might be due to the effects of invertase during ripening process²⁰; In another study, it became clear that raw fruits have variable amount of sucrose but during the process of ripening, sucrose cleaves to fructose and glucose.²¹ Sugars can also be seen in combination with flavonoids; derivatives of isoharmentin, like isoharmentin 3-β-D galactopyranoside that are present in the fruits.²¹ Among reducing sugars, xylose, mannose and rhamnose are also present in fruit extracts.¹³ To sum up, in every Russian olive trees, fructose and glucose seemed to be the major composition of the sugars but the amount of sugars does not depend on the origin or age of the trees.²¹

Alkaloids

Alkaloids were detected by TLC of different fractions of Russian olive. They are mainly condensed in the root, bark and aerial part of the plant. The most famous alkaloid of Russian olive is elaeagnin or calligonin which is structurally a tetrahydroharman. In the bark of this plant, N-methyl harmol, N-methyl tetrahydroharmol, Harman, dihydroharman, 2-methyl-1,2,3,4- tetrahydro-β-carboline, harmin and harmol are available as well.

There are some unlabeled medical usages of elaeagnin and β-carbolinesuch as the reduction of blood pressure and antimalarial effect, but none of them were investigated for the extracts of this plant.^{12,22-24}

Other ingredients

Vitamins and minerals play a major role in the medical properties of *E. angustifolia*. Vitamin A and K are oleophilic vitamins available in the methanol extract of the flowers of the plant. B vitamins are also present in the flowers.²⁵

The highest concentration of a mineral element in this plant belongs to potassium with a concentration of 8504 mg/kg. Second and third highest levels belong to sodium and phosphorus with 1731 and 635 mg/kg, respectively.²⁶ Calcium is also found with high levels in this plant and traditionally the flowers of Russian olive were considered a good source of Ca²⁺ among people of Iran.²⁵

Steroidal glycosides (saponins) can be detected in the fruits by Fontan-Candel method (foam-forming reaction). The saponins of Russian olive do not cause hemolysis of erythrocytes in isotonic saline. Aglycone part of

saponins can be either neutral or acidic. Total amount of saponins in the fruits of EA estimated to be 1.96 % of the net weight of the fruit.¹³

Free acids are also present in the fruits of this plant: benzoic acid, 4-hydroxy benzoic acid and vanilic acid are the benzoic derivatives. The derivatives of cinamic acid, like caffeic acid and ferulic acid are available as well. In this list, 4-hydroxy benzoic acid and caffeic acid are found in highest concentration, respectively.²⁰

Amino acids such as aspartic acid, threonine, serine, glutamine, proline, glycine, alanine, valine and some others were observed as well (Table 1).¹³

Traditional remedies

Traditionally, Russian olive was used as an anti-ulcer remedy for wound healing or sometimes gastric disorders.²⁷ *E. angustifolia* fruits were also famous in Turkish folklore as tonic, antipyretic, kidney disorder healing (anti-inflammatory and/or kidney stone treatment) and anti-diarrhea (astringent).^{16,28}

In ancient Iran, the fruit decoction of Russian olive was taught to be used a good remedy for fever, jaundice, asthma, tetanus and rheumatoid arthritis by Iranian apothecaries. In general, it was used as a substitute of any anti-inflammatory and analgesic agent, in the first line.²⁹

The leaves and fruits of the plant were also famous as diuretics and antipyretic agents. In Turkey, it was common to eat the fruits an hour before the meal as an appetizer.²¹ In Table 2 the traditional uses of this plant summarized. In the next sections, we will discuss some parts in detail.

Experimental and clinical studies

Anti-inflammatory and analgesic effects

Animal studies have shown the effectiveness of the aqueous and ethanol extract of Russian olive in pain and inflammation treatment. Flavonoids play the main role in this matter,³⁷ although anthocyanins, saponins³⁸ and terpenoids may take part as well.³⁹ There is a controversy on the success of the extract of this plant on acute or chronic pain. Formalin test is one of the reliable tests in both acute and chronic pain. In the first phase of formalin test, 5 minutes after injection, the direct influence of the compound on the pain fibers is tested and mostly called as an acute phase. 20 to 30 minutes after injection, the chronic phase or the inflammatory pain starts. For some medicines such

Table 1. Percentages of the active ingredients of dried ripe fruits of *Elaeagnus angustifolia*¹³

<i>Elaeagnus</i> spp. is in the family riparian trees growing near the rivers or water corridors	Percentage in dried ripe fruits of <i>E. angustifolia</i>
Reducing sugars	50.67-55.75%
Total sugar	60 ± 5%
Pectic polysaccharides	3.58 ± 0.3%
Total flavonoids and polycarboxilic acids	1.35 ± 0.15%
Total saponins	1.96 ± 0.52%
Ascorbic acid	5.6 mg%
B-carotene	17.5 mg%
Tannin	5.03 ± 0.05%

Table 2. Traditional remedies of *Elaeagnus angustifolia*

Extract/Country	Application	Rout of administration
Iran		
Infusion of the dried flowers	Anti-pyretic	oral ³⁰
Infusion of the dried leaves	Astringent	oral ³⁰
Turkey		
Fresh branch in hot water	To remove wart	External ³¹
Decoction of barks	Gall bladder problems and kidney stones	Drink one teacup twice a day for 20 days ³²
Extract of leaves with the ash of <i>Salix</i> sp. bark	To treat abscess	external ³³
Fruits	Kidney stones	oral ³⁴
Tea or decoction of root barks	Dysuria	Oral ³⁵
Extract of leaves with <i>Juglansregia</i> and <i>Menthalongifolia</i>	Sun block	External ³⁶
Azerbaijan		
Infusion of the aerobic parts	Treatment of alimentary canal	Oral (NAPALERT)

as morphine and codeine with a central effect both phases can be suppressed but for some others like non steroid anti-inflammatory drugs (NSAIDs) and steroids only the chronic phase can be effected.^{40,41}

While Farahbakhsh *et al.*³⁹ believe in ineffectiveness of the aqueous extract of the Russian olive in the acute phase; Hosseinzadeh *et al.*¹⁸ and Ahmadiani *et al.*²⁹ proved ($p < 0.001$) the comparable effect of the fruit extract with both NSAIDs and narcotics. Altogether, it can be inferred that the fruit extract of Russian olive with low doses (20-40 mg/kg) only suppresses the chronic pain²⁹ and high doses (130-450 mg/kg) can be successfully used against both chronic and acute pain.^{37,38}

On the other hand, the mechanism of its analgesic effects was also tested in hot-plate and writhing test. In comparison to indomethacin with a proven peripheral mechanism of pain relief in writhing test, the flavonoid extract of the Russian olive (134-402 mg/kg) showed both peripheral and central effects in hot-plate and writhing tests.³⁸

Quercetin is one of the most famous flavonoids with an analgesic effect available in Russian olive. It can block the activity of cyclooxygenase (COX) and lipoxygenase. *In vitro*, it has an inhibitory effect on the secretion of the immunoglobulins.¹⁰ Bioflavonoids with an ability to block the release of the bradykinin and arachidonic acid play an important role in the mechanism of controlling the chronic pain. Sometimes, muscle relaxant ability of the plant is known as an axillary mechanism for pain relief. Available alkaloids, such as harman and harmaline, inhibit mono amino oxidase (MAO) and increase serotonin and nor-adrenaline (like tricyclic antidepressants) at the synaptic sites. This mechanism can also be a part of analgesic effect of the plant.³⁷ Based on pharmacological information, flavonoids can block N-Methyl-D-aspartate (NMDA) receptor and reduce the amount of intracellular calcium and lead to reduced enzymatic activity of nitric oxide and phospholipase A2-Calcium-dependant protein.⁴² Flavonoids fill the ATP binding sites of kinases (serine/threonine or tyrosine kinase). These results obtained in less activation of phospholipase A2 and less production of COX. Some other flavonoids can block cAMP phosphodiesterase in the platelets. The reduction of cAMP chang-

es cytoskeletal rearrangements and deactivates special protein kinases. The substrates of the kinases like vasodilator-stimulated phosphoproteins remain inactive and finally platelet secretion, adhesion and aggregation stops. This is one of the important mechanisms in controlling inflammation.⁴³

Linoleic acid can also suppress the gene expression inflammatory cytokines. The inflammation is mediated by inhibition of NF- κ B (a protein complex that controls cytokine production) activation through PPAR-gamma pathway.⁴⁴

Recently in a clinical study, Russian olive extract (medulla powder and whole fruit, 15 g/kg, 8 weeks) showed a significant improvement in pain and inflammation management of knee osteoarthritis in women. It can reduce serum levels of inflammatory cytokines (TNF- α , MMP-1, etc) and also enhance patients' presentation of the disease; but the study was only carried out in obese women (according to reported body mass index) having chronic arthritis pain. It may need another comprehensive study containing at least pregnant women or athletes bearing acute pain and a range of dosages corresponding to previous laboratory studies.⁴⁵

Two separated randomized, double blind clinical studies, effectiveness of aqueous extract of Russian olive (300-600 mg/kg) which contained at least 0.21% kaempferol, was compared with ibuprofen (800 mg/kg) in women with knee osteoarthritis. *E. angustifolia* extract was very safe and tolerable in 2 doses and was so beneficial in reducing symptoms of osteoarthritis.^{46,47} In Table 3 most of the available studies are summarized. What we can infer is that analgesic effects of *E. angustifolia* are the most important medical applications of this plant. It is concentrated mainly in fruits and specially seeds of the trees. The mechanism does not relate to the opioid receptors, but the polyphenolic extracts can attenuates pains as good as the opioids.

Gastroprotective effect

From ancient remedies, Russian olive is famous for its antiulcer effect. *In vivo* studies in rats proved that the alcoholic extracts of this plant in low doses (400-800 mg/

Table 3. Anti-inflammatory and analgesic effects of *E. angustifolia*

Extract and doses	Compared medicine and doses	Experiment	Results and considerations	References
Seed (aqueous, ethanol and polyphenolic fraction) 100-1000 mg/kg	Indomethacin (1 mg/kg) Morphine (5 mg/kg)	Hotplate test Writhing test	Aqueous and ethanolic extracts were effective in concentration above 500 mg/kg Polyphenolic extract was effective in 300 mg/kg The role of polyphenolic groups in pain relief were proved	38
Total fruit (aqueous fraction) 1000 mg/kg	Sodium salicylate (100-300 mg/kg)	Formalin test Tail-flick test Anti-inflammatory test	Acute phase of pain: peritoneal injection was effective Chronic phase of pain: like NSAIDs have pain relief effects even orally In both phases, naloxone could not reverse the effects Central mechanism was proved	29
Total fruit (aqueous fraction) 50 mg/kg	Morphine (10 mg/kg) Indomethacin (10 mg/kg) Dexamethasone (10 mg/kg)	Formalin test	No suppression in acute phase (because of low doses of extract) Pain relief in chronic phase with an inhibition of COX enzymes naloxone couldn't reverse the effects	39
Fruit (aqueous fraction) 20 mg/kg	Diclofenac (10 mg/kg)	Xylene-induced ear edema	Seeds have the most anti-inflammatory effect in chronic phase	18
Fruit (aqueous fraction) 250-500-700 mg/kg	Indomethacin (5 mg/kg)	Writhing test	Comparable visceral pain relief with NSAIDs	37
Fruit (aqueous fraction) 1-1.5 g/kg	Imipramine (40 mg/kg)	Hot-plate test	Comparable pain relief in chronic phase with imipramine	37

kg) and high doses (3150 mg/kg) can improve lesions of NSAIDs and ethanol, respectively. These properties are as effective as misoprostol and less operative than omeprazole. In histopathological examinations, no protective effect on stomach mucosa has been observed. In another study, aqueous extract of Russian olive imposed positive effects on pregnant mice. With doses as low as 640 mg/kg, fetal stomach cells became hypertrophic by gene regulation. In other words, *E. angustifolia* extract could reinforce fetus digestive system. There are also some evidences that Russian olive fruit extract can reduce gastric acid secretion simulated by cholinergic system in a dose and time dependent manner. Russian olive was shown to relax the muscle contraction of lumen and with its antioxidant effects; it works as a barrier on the surface of stomach against gastric mucosa.^{28,48-50}

We can infer from this section that Russian olive ethanol extract is comparable with misoprostol on gastric ulcer lesions; it can reduce mucosal damage formed by aspirin, NSAIDs and alcohol like H₂ receptor blockers, but not as powerful as proton-pump inhibitors. With less side effects (especially no oxytocic property), Russian olive dosage forms or extracts can be administered before NSAIDs-intake or for pregnant women in controlled quantities.⁵¹

Wound treatment

In folk remedies, the extracts of Russian olive are known as a wound healing accelerator. If treatment of inflammation and proliferation is considered as the main steps of wound treatment, Russian olive fruit extracts can help in wound closure. It can increase the hydroxyproline content, improve the histological scores (epidermis regener-

ation, collagen deposition and proliferation), control the pain and inflammation and inhibit the cyclooxygenases I and II. Moreover, *E. Angustifolia* has a potent antibacterial property comparable with mupirocin 2% with added antioxidant activity. Phenolic compounds, mostly flavonoids are responsible for these effects. Vitamin A also plays important roles; the acceleration of cell replication, more collagen precipitation and induction of hyaluronate synthesis are the main effects of this vitamin in wound healing.^{39,52-55}

Antibacterial and antifungal effect

In general, this plant shows lower overall antimicrobial effects in comparison to other similar species.⁵⁶ However, in some cases, extraordinary effects have been reported. Reported antifungal and antibacterial effects vary based on the dose used and the extraction method recruited. In an in vitro study, its leaf extract was used to treat mastitis pathogens.⁵⁷ These pathogens cause severe illnesses in domestic animals and by triggering steep decline in milk production result in vast economical loss.⁵⁸ Such illnesses are also classified as threatening to public human health.⁵⁹ They are caused by a group of bacteria such as *Staphylococcus aureus* and Coagulase Negative *Staphylococci* (CNS) which are susceptible to this plant's extracts.⁶⁰ In another study, four hospital germs were collected and tested in a disc diffusion method, and two soft extracts of leaves and flowers of Russian olive proved to have an intense antibacterial activity against wound germs (*S. aureus*), pharynx exudates (*Streptococcus pyogenes*), saliva (*Klebsiella pneumoniae*) and urine (*Escherichia coli*).⁶¹ A brief review of antibacterial and antifungal activity of Russian olive is gathered in Table 4. In accordance with the

Table 4. Different extracts of *Elaeagnus angustifolia* and their effects on pathogens

	Aqueous	Ethanollic	Chloroform	Crude	n-Hexane	Methanolic
<i>Mastitis pathogen</i> ⁶⁰						√
<i>Staphylococcus aureus</i> ⁶²		√				
<i>Escherichia coli</i> ⁶²	√					
<i>Salmonella thyphimurium</i> ⁵⁷	√	√			√ (The most effective extract)	√
<i>Bascillusthuriensis</i> ⁵⁶		√				
<i>Pseudomonas aeruginosa</i> ⁶²		√				
<i>Pseudomonas extorquous</i> ⁵⁶		√				
<i>Yersinia enterocolitica</i> ⁶³						√
<i>Candida albicans</i> ⁵⁷	√	√			√ (The most effective extract)	
<i>Aspergillus fumagatus</i> ⁶²				√	√ (The most effective extract)	
<i>Aspergillus favis</i> ⁶²	√	√			√ (The most effective extract)	
<i>Aspergillus niger</i> ⁶²		√			√ (The most effective extract)	
<i>Alternariasolan</i> ⁶⁴						√
<i>Bacillus subtilis</i> ⁶⁵	√	√			√	
<i>Shigelladysentriae</i> ⁶⁵	√	√			√	
<i>Vibrio cholerae</i> ⁶⁵	√	√			√	

importance of this plant in the traditional remedies, we considered some of its applications were related to its fungicidal and bactericidal properties. In ancient apothecary, it was said that *E. angustifolia* was administered to control the gull bladder problems and diarrheas. Gull bladder problems may be *E.coli* related or the table-listed gram negative bacteria could be responsible for the mentioned diarrhea and so on.

The exact mechanism of these properties is not clear, but most scientists believe in phenolic compounds,^{52,62} alkaloids⁶² and essential oils⁶⁰ as predominant reasons of being germicide in medical plants. Amongst its flavonoids, flavones like kaempferol, quercetin and isohamentin are deemed to provide such pharmacologic effects.⁵² Flavonoids especially the ones with more lipophilic structures can penetrate into the cell wall of the bacteria easily and spoil its condensed structure. The intracellular components permeate and the bacteria are being killed. Another proposed mechanism is related to the complex of the flavonoids with active proteins of the bacteria with specific (covalent) or nonspecific (hydrogen or van der Waals) bonds and make critical changes in bacteria's function that may finally lead to the death of the microorganism.⁴³

Antioxidant effect

Most studies on *E. angustifolia* is related to its antioxidant capacity of phenolic compounds and antocyanosides. The highest content of total phenolic compounds can be extracted on the first 10 days of October, therefor the best time for harvesting Russian olive is exactly at this time in cases highest antioxidant properties are desired.^{66,67} Apart from the seasonal changes, genotype is also important in the amount of antioxidant capacity. For *E. angustifolia* we know at least 7 genotypes (IEa1 to 7) in Iran.⁶⁸

Polyphenols can suppress the oxidative stress produced by reactive oxygen species (ROS) with their hydroxyl groups. They can also chelate metals.⁶⁹ Antioxidant activity of raw

materials can be mainly evaluated in two methods: antioxidant capacity against of free radical species comprising of hydrogen atoms transfer reactions model (HAT), single electron transfer reactions model (SET) and a hydrogen-electron transfer models which is a mixture of HAT and SET methods; and antioxidant capacity against biological markers and characterized substrates.⁷⁰

Oxygen radical absorbance capacity assay (ORAC), a HAT mechanism, is a core method in antioxidant studies. The senior method refers to the usage of β -phycoerythrin, a fluorescent hydrophilic probe, which was photo sensitive and caused false positive results. Fluorescein, eosin and 6-carboxyfluorescein with higher stabilities and more specific interactions with antioxidants are the probes of ORAC method.⁷¹ Phenolic compounds and even vitamins of Russian olive can hydrogenate peroxy radicals of AAPH and increases the fluorescent emission strength. Among the extracts, water/ethanol (1:1) extracts of the leaves has the most potent antioxidant property. All in all, this method has exclusively been characterized to evaluate the antioxidant capacity of water-soluble phytochemicals and may not cover all aspects of this plant.^{72,73}

Furthermore, 2, 2-Diphenyl-1-picrylhydrazyl (DPPH) radical assay based on Benvenuti *et al.* method is an available and cheap antioxidant analysis experiment. The stable and commercially accessible DPPH is a free radical that can accept hydrogens and electrons from polyphenols and also is one of the comparable methods of antioxidant measurement (Fig. 3).^{70,74}

With an increase in the concentration and polarity of ethyl acetate extracts of Russian olive, its polyphenolic contents can clear DPPH radicals out up to 94%. It has been estimated and shown that its antioxidant ability is comparable with α -tocopherol and BHT in terms of IC₅₀. Polysaccharide contents of Russian olive comprising of monomers such as rhamnase, xylose, mannose, glucose and galactose can reduce DPPH free radicals up to 88.1%. From the con-




Fig. 3. Anti-inflammatory mechanism of *E. angustifolia*: Flavonoids of *E. angustifolia* can directly block active sites of COX and the secretion of the arachidonic acid. Thus, arachidonic acid cannot change into prostaglandins and leukotrienes, the most important mediators of vasodilation, platelet activation and inflammation (filled arrows). Another proposed mechanism describes the importance of flavonoids in blocking NMDA receptors. NMDA receptor inactivation leads to a significant calcium secretion reduction from PLC pathway and finally no kinase, PLA2 and COX activation, respectively (dashed arrow). (N-methyl-D-aspartate: NMDA, phospholipase C: PLC, inositol trisphosphate: IP3, phospholipase A2: PLA2, cyclooxygenase: COX)

centration point of view, polyphenols radical scavenging activity is more than ten times stronger than polysaccharides.⁷⁵⁻⁷⁸ As a suggestion, it is better to define antioxidant ability of the raw materials with a standard chemical like Trolox instead of IC₅₀ or percentage. Trolox which can be the positive control in research projects and the minimum inhibitory concentration, can become a measure of comparison in chains of studies.⁶³ However, what we are sure about is that the antioxidant capacity of this plant has a linear correlation with the amount of phenolic compounds.⁷⁹ Phenolic compounds can be extracted from the methanol or ethanol extracts. They can donate hydrogen or electron and remove the free radicals. They reduce the amount of ROS by controlling the free-radical-producer enzymes like NADH oxidase, glutathione s-transferase and micro-

somal monooxygenase or chelating trace elements. Some of active ingredients of *E. angustifolia* such as epicatechin, catechin, rutin, quercetin and kaempferol with a hydroxyl group in seventh position of the 4-chromanone ring of the base structure have the maximum efficacy against oxidative groups.⁸⁰

Anti-neoplastic effect

With an increase in concentration and polarity of *E. angustifolia* extracts, its anti-tumor effects level up. In an *in vivo* study on mice, elevation in immunity with increase in spleen index and thymus index manifestation has been reported as the main anticancer mechanism of Russian olive extracts. It has been shown that the anti-cancer properties are the effects of essential oils (ethyl cinnamate, 2-phenyl-ethyl benzoate, 2-phenyl-ethyl isovalerate, nerolidole, squalene and acetaphenone), flavonoids and proanthocyanosides.^{75,81,82}

Moreover, the efficacy of Russian olive in inhibition of angiogenesis in Human umbilical vein endothelial cells may result in control of diabetes, retinopathy, rheumatoid arthritis and skin diseases like psoriasis in addition to malignancies.^{83,84} Documented studies on Russian olive extracts in anti-neoplastic effects have been summarized in Table 5.

Flavonoids can down regulate mutant p53 genes (p53 wild gene is a regulatory gene that codes special proteins which control cell cycles and known as a tumor suppressor gene) and arrest the cell cycle in G2 or M in malignant tumors. At the same time, flavonoids can suppress the expression of Ras protein and control the heat shock proteins especially in leukemia and colorectal cancers. Quercetin, one of the main flavonoid compositions of *E. angustifolia* is known as an anti-proliferative agent because it can block the cell surface tyrosine kinases and stops the transfer of growth messages to the nucleus.⁴³

Muscle relaxant

In traditional medicine, muscle relaxant effects of EA has been noted upon in different literature.¹⁶ An *in vivo* study on mice to investigate such effects showed that polar extracts of this plant can exhibit dose dependent effects comparable to diazepam (2 mg/kg). It was suggested that flavones available in such extracts possess a partial agonistic effect on benzodiazepine (BDZ) receptors.⁸⁸ Flavones

Table 5. Anti-neoplastic effects of *Eaeagnus angustifolia*

Type of extract	Cell line	Considerations	References
Hydroalcoholic of flowers	Human umbilical vein endothelial cells(HUVEC)	<i>In vitro</i> inhibition of angiogenesis	84
Phenolic extract	Human colorectal adenocarcinoma cells(HT29)	<i>In vitro</i> control of colorectal cancers in different stages	85, 86
Ethanollic of leaves	Breast adenocarcinoma cell(MCF7)	<i>In vitro</i> studies showed no significant difference with control group	87
Ethyl acetate of the whole plant	Human cervical carcinoma cells (Hela)	<i>In vitro</i> study	75
Edible parts of the plants	Human hepatoma cells(HepG2)	<i>In vitro</i> and <i>in vivo</i> studies on H22 cells-bearing mice	81
Tincture of flowers	Walker 256 carcinoma cells	<i>In vivo</i> study on Wistar rats	69

especially the 6-substitution compounds have strong affinity with GABA_A receptors; however flavones which are naturally synthesized, have partial agonistic effect on BDZ receptors. This mechanism is mainly described for the sedative compounds; but in this filed, the extract shows a pharmacological effect comparing to diazepam for controlling muscle contractions. After the activation of GABA receptors, chloride channels open and hyperpolarize cell membrane which prevents further excitations.⁸⁹ On the other hand, the same muscle relaxant effect is produced in smooth muscles of the intestine, due to the acetylcholine antagonizing effects of the said extract. This provides a means to treat intestinal colic pains.⁹⁰

Cardiovascular effects

Elaeagnin, one of the alkaloids of *E. angustifolia*, was reported as a blood pressure optimizer. It has a tetrahydroharman structure resembling reserpine on the basis of overlay studies of bioinformatics (Fig. 4). It can bind reversibly or irreversibly to the human monoamine oxidase A (MAO-A) active site and lower the blood pressure. As elaeagnin can fit with X-ray structure of harmine in MAO-A active site, theoretically, it would be effective as a blood pressure controller like reserpine (harmine in white and elaeagnin in yellow). On the other hand, aqueous leaf extracts of *E. angustifolia* with antioxidant property, can improve oxidative stress state brought by ischemic/reperfusion (I/R) in rats. This extract (0.5 mg/mL and 1 mg/mL) can reduce cardio toxicity of I/R and increase myocardial biochemical parameters. Altogether, this plant with possible reduction of blood pressure and enhancing the recovery of I/R can be a good choice in cardiac diseases but more studies needed.^{37,91} ROS production is one of the main mechanisms of myocardial dysfunction and necrosis after I/R. superoxide dismutase (SOD) enzymes can control the radical species or ROS in myocardium. But, the problem triggers when the amount of ROS exceed the threshold limit of SOD. The leaves extract of *E. angustifolia* with polyphenolic compositions can reduce the activity of ROS, serum level of carbonyl groups and malondialdehydes with the molecular mechanism described in Fig. 5 and increase the ability of SOD to control the oxidative stress.⁹²




Fig. 4. Overlay of elaeagnin (the active component of *E. angustifolia*) and harmine

Other medical applications

In animal models, aqueous extract of *E. angustifolia* can improve cognitive disorders. Tamtaji *et al.* treated the scopolamine-induced alzheimeric rats with increased doses (50-100-200-400 mg/kg) of Russian olive and resulted in significant improvements of special learning and memory.⁹³

In a randomized clinical trial, females with orgasmic disorders received flower extract of *E. angustifolia* (4.5 g/D in 2 doses) and sildenafil citrate (50 mg, 1 h before intercourse), female sexual function and the levels of prolactin and TSH was measured after 4 weeks' treatment. Hormonal levels were found to remain even. Although Russian olive was less functional than sildenafil, it was effective in reduction of the frequencies of orgasmic disorders. The mechanism is not clear, but it may be the result of NO rise and subsequently cGMP increase in cells. It may cause vaginal smooth muscle relaxation, artery vasodilation and finally swelling of genital system.⁹⁴

The effects of Russian olive in treatment of gastrointestinal problems have been proved. It may also be operative in ulcerative colitis and help to prevent cancers. Histopathological studies in rats with induced ulcerative colitis presented a significant improvement with edible extract (600 mg) and less improvement with the enema gel containing 20 percent extracts of Russian olive.⁹⁵




Fig. 5. Free radical scavenging mechanism of flavonoids.

Drug formulation

Using herbal compounds as drug carriers due to their lower side effects is a good step forward.⁹⁶ In this category, several studies have been carried out using dried extract to transport calcium carbonate, naproxen and ibuprofen trying to increase drug wetting, prevent caking, increase particle surface and increase drug dissolution speed. These efforts are expected to result in increased blood concentration of medicines. Nanocapsules containing calcium carbonate and EA in addition to other formulation improvements in comparison to calcium carbonate nanocapsules and calcium carbonate tablets result in higher calcium blood levels. Solid dispersions containing naproxen and ibuprofen show higher analgesic and anti-inflammatory effects. Despite its lower efficacy, in comparison to chemicals such as cross povidone, extracts of this plant are still more favorable, in large part due to less documented side effects.⁹⁷⁻⁹⁹

Moreover, 6% soft extract of Russian olive can be useful in cream formulations. In addition to the antioxidant and anti-inflammatory effects for wound healing, it helps the spread capacity and thixotropic properties of cream or ointments bases. The best formulation for the extract insertion is an anionic base (oil/water emulsions).^{69,100,101}

Russian olive extracts have also been inserted in topical gels for oral lesions. Clinical studies on 28 patients with oral lichen planus resulted in considerable decrease of lesion size and pain. Minerals and vitamins especially vitamin K with its efficacy in blood coagulation play main rolls in this matter.¹⁰²

Use in pregnancy controversies

Previously, in traditional benefits of Russian olive, it was described that its extract can be useful for the treatment of rheumatoid arthritis. Pregnant women were among the suggested groups by the alchemists to use this extract against pain, inflammation and all problems in their joints and bones. Recent studies have shown the disadvantages of the fruit extract on the embryo osteogenesis and chondrogenesis. Although this extract does not reduce the amount of calcium, it reduces the volume of bones and cartilages. In contrary, mouse embryos showed a significant increase in femur length. The exact mechanism is not clear, but we can say phytoestrogens like flavonoids and tannins may inhibit non-classic estrogen signaling pathway, G protein receptor 30 (GRP30), and exert these effects in a dose depending manner. To end up, Russian olive extract can improve osteogenesis and chondrogenesis, but may decrease bone mass in mouse embryos, although it makes no craniofacial malformation or limb abnormalities.⁴⁸ Therefore, in this case, pregnant mothers should avoid the traditional remedies for joint and bone disorders.¹⁰³

Possible adverse effects

Russian olive is among the safest trees in the world and apart from possible allergies, there is no other hazard reports. Even though it is rich in heavy metals such as Cr,

Pb, Zn, Cu, Ni and Co, some researches proved that in any condition of harvesting, the amount of heavy metals in all parts of the tree will not exceed the WHO limits (based on WHO guideline: the permissible limit of Cr, Pb, Zn, Cu, Ni and Co in medicinal plants is 1.5, 10, 50, 10, 1.5 and 0.2 part per million, respectively).^{62,104} Even in an investigation on pregnant mice, *E. angustifolia* extracts made no significant differences in weight and CRL of mice fetus; but, because of the effects of concentrated extracts on bones and cartridge volumes, especially in pregnant women, more consideration may be needed.⁴⁸

Allergenicity

Russian olive is common as a wind-cutting tree in American and European countries, so its pollens would spread widely in the cities especially in the pollination season, spring. In an investigation, Ole e 1 and Ole e 4-like allergens, major allergens in olive pollen,^{105,106} were recognized in this plant. With a prick test, 30 % of patients with rhinitis and rhinoconjunctivitis were sensitive to Russian olive pollens and 44 percent of patients with sensitivity to pollens have positive Prick tests. With these findings and the fact that its pollen size is much greater than grass and olive pollens (40-50 µm), it can be concluded that *E. angustifolia* is one of the major causes of allergies in the cities.¹⁰⁷ The mechanism which triggers these reactions is an IgE-mediated type II allergic type.¹⁰⁸

Moreover, dermatitis caused by contact with *Elaeagnus* sp. has already been reported; but, mostly among florists with repeated contact.¹⁰⁹

Concluding remarks

Utilizing the unlimited natural sources of medicines with the least restrictions of safety is one of the human beings goals. In this area, *E. angustifolia* can fulfill these objectives. This plant is a riparian tree considered as a N2-fixer herb, increases the nitrification of the soil (inorganic nitrogen).¹¹⁰ It is rich in water and fat soluble vitamins, flavonoids, carbohydrates, alkaloids and biological active lipids. This review aimed to gather ethnopharmacological properties of *E. angustifolia* with focus on active constituents and medical benefits, and also make suggestions for future plans. For this purpose, almost all the medical articles from 1970 to 2016 were collected from various databases such as Embase, NCBI, Science Direct and Scopus. The most important and peer-reviewed information were analyzed and classified judiciously.

For years, *E. angustifolia* was used as an astringent, kidney stone removal, anti-inflammatory and pain relief agent. It was a well-known remedy for fever, jaundice, asthma, tetanus and rheumatoid arthritis. In turkey, *E. angustifolia* was an appetizer and a source of nutrition. It can be a food additive; as in an experiment by Cakmakc *et al.* addition of the flour and crust of the fruits as a flavor and formulation optimizer (viscosity enhancement) in an ice cream have showed perfect results. In food industry, natural plants with antioxidant properties can result in vast commercial advertisements.¹¹¹

Effects of Russian olive on inflammation and pain have been studied the most. More than 4 clinical trials are available and all of them have reported a comparable efficacy as NSAIDs. Total fruit has muscle relaxant capacity in a dose-dependent manner comparable to that of diazepam, COX inhibition property like indomethacin and ibuprofen, and is full of kaempferol that can suppress inflammation mediators.

A number of experiments have evaluated its active constituents for their applications in malignancies, ulcers, gastric problems and osteoarthritis. Previous studies showed that even in injection forms in mice and rats, there is no significant immunological responses (as the corticosteroid levels were even),³⁹ this plant can even be applied in developed injection formulations for special applications like intra-articular vials.

In “experimental and clinical studies” section, the *E. angustifolia* extract was suggested as blood pressure optimizer, because of special structure similarity of elaeagnin to hamine and reserpine. Till now, no supported studies have proved it; perhaps due to the time-consuming mechanism of reserpine in controlling blood pressure. As known, reserpine blocks MAO irreversibly and then increases the amount of neurotransmitters in the synaptic cavity; this elevation of neurotransmitters lead to down regulation of the vascular receptors and decreases blood pressure subsequently. The aforementioned process is a long-lasting effect. None of the clinical studies of *E. angustifolia* extract consider time as a major factor in cardiovascular protective properties of the plant.¹¹²

Flavonoids of this plant are so hydrophobic. Lots of *E. angustifolia* medicinal benefits are related to this property of flavonoids. Anti-bacterial effects are one of them. Flavonoids can penetrate the cell membrane of the bacteria and exert their destructive effects. In most cases, aggregation of bacteria was seen. Some of its flavonoids like kaempferol and quercetin show slight β -lactamase inhibitory effects, as well. As presented in Table 4, the main antibacterial properties are related to the methanol or ethanol extracts.¹¹³

The lack of evidence on the relation of the medical applications and detailed structure of constituents, conceal its exclusive properties. There are no reports on experimental studies of pharmacologic/toxicologic effects of a few compounds like elaeagnine or elaeagnoside that are found in this plant. The gastroprotective effects of the plant appear to lead us towards development of medicines comparable with misoprostol. As for the anti-neoplastic effects or bactericidal effects, this plant can be a source of inspiration for rational drug design and medicinal chemistry. As a matter of fact, around 85 percent of chemotherapy agents have natural sources.¹¹⁴ However, the lack of clinical studies on this plant has limited its availability in drug forms. It is certain that further studies would be necessary to fill such information paucity and gap.

Ethical approval

There is none to be declared.

Review Highlights

What is current knowledge?

✓ *E. angustifolia* can be a source of bioflavonoids that are so applicable in controlling pain and inflammation.

✓ Active ingredients of *E. angustifolia* can even be useful as anti-cancer, anti-bacterial and anti-fungal medicines.

✓ The protective properties of *E. angustifolia* extract in gastrointestinal system became clear.

What is new here?

✓ The relationship between its active ingredients and medical properties are not mechanistically clear; but there are some evidences in participation of the flavonoids of Russian olive in controlling enzymes (COX) or blocking the receptors (NMDA-R or BDZ-R).

✓ One of the possible mechanisms in which active ingredients of Russian olive may decrease the blood pressure, have been discussed.

✓ No critical and significant side effects of this plant (even in high doses) have been reported.

Competing interests

There is none to be declared.

Acknowledgments

This review article was supported by research center of Mashhad University of Medical Sciences.

References

1. Mineau MM, Baxter GV, Marcarelli AM, Minshall GW. An invasive riparian tree reduces stream ecosystem efficiency via a recalcitrant organic matter subsidy. *Ecology* **2012**;93:1501-8.
2. Lamers JP, Khamzina A. Seasonal quality profile and production of foliage from trees grown on degraded cropland in arid Uzbekistan, Central Asia. *J Anim Physiol Anim Nutr (Berl)* **2010**;94:e77-85. doi: 10.1111/j.1439-0396.2009.00983.x.
3. Mehrabani Natanzi M, Pasalar P, Kamalinejad M, Dehpour AR, Tavangar SM, Sharifi R, *et al*. Effect of aqueous extract of *Elaeagnus angustifolia* fruit on experimental cutaneous wound healing in rats. *Acta Med Iran* **2012**;50:589-96.
4. Kurdali F, Al-Shamma'a M. Natural abundances of ¹⁵N and ¹³C in leaves of some N₂-fixing and non-N₂-fixing trees and shrubs in Syria. *Isotopes Environ Health Stud* **2009**;45:198-207. doi: 10.1080/10256010903084126.
5. Huang Z, Liu M, Chen B, Uriankhai T, Xu C, Zhang M. Distribution and interspecific correlation of root biomass density in an arid *Elaeagnus angustifolia*-*Achnatherum splendens* community. *Acta Ecol Sin* **2010**;30:45-9. doi: 10.1016/j.chnaes.2009.12.008.
6. Gonzalez-Munoz N, Castro-Diez P, Fierro-Brunnenmeister N. Establishment success of coexisting native and exotic trees under an experimental gradient of irradiance and soil moisture. *Environ Manage* **2011**;48:764-73. doi: 10.1007/s00267-011-9731-3.
7. Rood SB, Braatne JH, Goater LA. Favorable fragmentation: river reservoirs can impede downstream expansion of riparian weeds. *Ecol Appl* **2010**;20:1664-77.
8. Fritz Hans Schweingruber AB, Ernst-Detlef Schulze. *Elaeagnaceae. Atlas of Stem Anatomy in Herbs, Shrubs and Trees*. New York: Springer; 2011. p. 152-5.
9. Farzaei MH, Bahramsoltani R, Abbasbadi Z, Rahimi R. A comprehensive review on phytochemical and pharmacological aspects of *Elaeagnus angustifolia* L. *J Pharm Pharmacol* **2015**;67:1467-80.
10. Kumar R, Kaur M, Silakari O. Chemistry and biological activities

- of thioacridines/thioacridones. *Mini Rev Med Chem* **2013**;13:1220-30.
11. Wang Y, Guo T, Li JY, Zhou SZ, Zhao P, Fan MT. Four flavonoid glycosides from the pulps of *Elaeagnus angustifolia* and their antioxidant activities. In: Cai SZ, Zhang QF, eds. *Advanced Materials Research*. Nanjing, Jiangsu; **2013**. p. 16-20.
 12. Bekker NP, Glushenkova AI. Components of Certain Species of the Elaeagnaceae Family. *Chem Nat Compd* **2001**;37:97-116. doi: 10.1023/a:1012395332284.
 13. Abizov EA, Tolkachev ON, Mal' Tsev SD, Abizova EV. Composition of biologically active substances isolated from the fruits of Russian olive (*Elaeagnus angustifolia*) introduced in the European part of Russia. *Pharm Chem J* **2008**;42:696-8.
 14. Kusova RD, Luk'yanchikov MS. Fatty acid composition of the fruit oil of *Elaeagnus angustifolia*. *Chem Nat Compd* **1989**;25:718. doi: 10.1007/bf00598274.
 15. Goncharova NP, Plugar VN, Rashkes YV, Isamukhamedov AS, Glushenkova AI. Oxygenated fatty acids of the seeds of *Elaeagnus angustifolia*. *Chem Nat Compd* **1994**;30:661-5. doi: 10.1007/bf00630597.
 16. Goncharova NP, Glushenkova AI. Lipids of elaeagnus fruit. *Chem Nat Compd* **1990**;26:12-5. doi: 10.1007/bf00605188.
 17. Zhang N, Bao D, Yang C, Huang X. Ultrasound-assisted extraction of *Elaeagnus angustifolia* seeds oil and its physical-chemical properties. *Journal of the Chinese Cereals and Oils Association* **2013**;28:82-5.
 18. Hosseinzadeh H, Rahimi R. Anti-inflammatory effects of *Elaeagnus angustifolia* L. fruits in mice and rats. *Iran J Basic Med Sci* **1999**;24:143-7.
 19. Ayaz M, Riaz M, Malik A, Ahmad E, Fatima I, Arif Lodhi M, et al. Elaeagnoside, chymotrypsin inhibiting steroidal glucoside from *Elaeagnus orientalis*. *Nat Prod Res* **2009**;23:409-14. doi: 10.1080/14786410601083506.
 20. Ayaz FA, Bertoft E. Sugar and phenolic acid composition of stored commercial oleaster fruits. *J Food Compos Anal* **2001**;14:505-11. doi: 10.1006/jfca.2001.1004.
 21. AYAZ FA. *Soluble Sugar Composition of Elaeagnus angustifolia* L. var. *orientalis* (L.) Kuntze (Russian olive) Fruits. *Tr J Bot* **1999**;23:349-354.
 22. Nikolaeva AG. Alkaloids of *Elaeagnus angustifolia*. *Chem Nat Compd* **1970**;6:659. doi: 10.1007/bf00563479.
 23. Allen JR, Holmstedt BR. The simple β -carboline alkaloids. *Phytochemistry* **1980**;19:1573-82.
 24. Ancolio C, Azas N, Mahiou V, Ollivier E, Di Giorgio C, Keita A, et al. Antimalarial activity of extracts and alkaloids isolated from six plants used in traditional medicine in Mali and Sao Tome. *Phytother Res* **2002**;16:646-9.
 25. Esmaili A, Niknam S. Characterization of nanocapsules containing *Elaeagnus angustifolia* L. extract prepared using an emulsion-diffusion process. *Flavour Fragr J* **2013**;28:309-15. doi: 10.1002/ffj.3164.
 26. Cansev A, Sahan Y, Celik G, Taskesen S, Ozbey H. Chemical Properties and Antioxidant Capacity of *Elaeagnus angustifolia* L. Fruits. *Asian J Chem* **2011**;23:2661-5.
 27. Yeşilada E, Gürbüz İ. A compilation of the studies on the Anti-ulcerogenic effects of medicinal plants. *Recent Progress in Medicinal Plants. Phytochem Pharmacol* **2002**;2:111-74.
 28. Gürbüz I, Üstün O, Yesilada E, Sezik E, Kutsal O. Anti-ulcerogenic activity of some plants used as folk remedy in Turkey. *J Ethnopharmacol* **2003**;88:93-7.
 29. Ahmadiani A, Hosseiny J, Semnianian S, Javan M, Saeedi F, Kamalinejad M, et al. Antinociceptive and anti-inflammatory effects of *Elaeagnus angustifolia* fruit extract. *J Ethnopharmacol* **2000**;72:287-92. doi: 10.1016/s0378-8741(00)00222-1.
 30. A. Z. *Medicinal plants*. 5th ed. Tehran, Iran: Tehran University Publications; **1992**.
 31. Sezik E, Yeşilada E, Honda G, Takaishi Y, Takeda Y, Tanaka T. Traditional medicine in Turkey X. Folk medicine in Central Anatolia. *J Ethnopharmacol* **2001**;75:95-115.
 32. Kültür S. Medicinal plants used in Ki{dotless}rklareli Province (Turkey). *J Ethnopharmacol* **2007**;111:341-64.
 33. Yesilada E, Honda G, Sezik E, Tabata M, Fujita T, Tanaka T, et al. Traditional medicine in Turkey. V. Folk medicine in the inner Taurus Mountains. *J Ethnopharmacol* **1995**;46:133-52.
 34. Honda G, Yesilada E, Tabata M, Sezik E, Fujita T, Takeda Y, et al. Traditional medicine in Turkey. VI. Folk medicine in west Anatolia:Afyon, Kutahya, Denizli, Mugla, Aydin provinces. *J Ethnopharmacol* **1996**;53:75-87. doi: 10.1016/S0378-8741(96)01426-2.
 35. Fujita T, Sezik E, Tabata M, Yesilada E, Honda G, Takeda Y, et al. Traditional medicine in Turkey VII. Folk medicine in middle and west Black Sea regions. *Econ Bot* **1995**;49:406-22.
 36. Sezik E, Yeşilada E, Tabata M, Honda G, Takaishi Y, Fujita T, et al. Traditional medicine in turkey viii. Folk medicine in east Anatolia; Erzurum, Erzincan, Ağrı, Kars, İğdir provinces. *Econ Bot* **1997**;51:195-211.
 37. Karimi G, Hosseinzadeh H, Rassoulzadeh M, Razavi BM, Taghiabadi E. Antinociceptive effect of *Elaeagnus angustifolia* fruits on sciatic nerve ligated mice. *Iran J Basic Med Sci* **2010**;13:97-101.
 38. Ramezani M, Hosseinzadeh H, Daneshmand N. Antinociceptive effect of *Elaeagnus angustifolia* fruit seeds in mice. *Fitoterapia* **2001**;72:255-62. doi: 10.1016/s0367-326x(00)00290-2.
 39. Farahbakhsh S, Arbabian S, Emami F, Moghadam BR, Ghoshooni H, Noroozadeh A, et al. Inhibition of cyclooxygenase type 1 and 2 enzyme by aqueous extract of *Elaeagnus Angustifolia* in mice. *Basic Clin Neurosci* **2011**;2:31-7.
 40. Abbadie C, Taylor BK, Peterson MA, Basbaum AI. Differential contribution of the two phases of the formalin test to the pattern of c-fos expression in the rat spinal cord:studies with remifentanyl and lidocaine. *Pain* **1997**;69:101-10.
 41. Hunskaar S, Hole K. The formalin test in mice:dissociation between inflammatory and non-inflammatory pain. *Pain* **1987**;30:103-14.
 42. Rang HP, Ritter JM. *Text Book of Pharmacology*. 7th ed. New York**2012**.
 43. Kumar S, Pandey AK. Chemistry and biological activities of flavonoids:an overview. *ScientificWorldJournal* **2013**;2013:162750. doi: 10.1155/2013/162750.
 44. Zhao G, Etherton TD, Martin KR, Vanden Heuvel JP, Gillies PJ, West SG, et al. Anti-inflammatory effects of polyunsaturated fatty acids in THP-1 cells. *Biochem Biophys Res Commun* **2005**;336:909-17. doi: 10.1016/j.bbrc.2005.08.204.
 45. Nikniaz Z, Ostadrahimi A, Mahdavi R, Ebrahimi AA, Nikniaz L. Effects of *Elaeagnus angustifolia* L. supplementation on serum levels of inflammatory cytokines and matrix metalloproteinases in females with knee osteoarthritis. *Complement Ther Med* **2014**;22:864-9.
 46. Ebrahimi A, Ebrahimi F, Nikniaz Z, Mahdavi R, Ostadrahimi A, Nikniaz L. Effect of elaeagnus angustifolia L. Medulla fruit powder on pain of females with knee osteoarthritis. *Int J Rheum Dis* **2015**;18:79-80.
 47. Panahi Y, Alishiri GH, Bayat N, Hosseini SM, Sahebkar A. Efficacy of *Elaeagnus angustifolia* extract in the treatment of knee osteoarthritis:A randomized controlled trial. *EXCLI J* **2016**;15:203-10.
 48. Malihezaman M, Mahbobe M, Maryam SL. The ultrastructural and stereological study of aqueous extracts of *Zataria multiflora* Boiss and *Elaeagnus angustifolia* on the mouse fetus stomach. *J Biol Sci* **2007**;7:648-52.
 49. Eliassi A, Ghodrati M, Kamalinrad M. Effect of oral (systemic) *Elaeagnus angustifolia* L. fruit on stimulated gastric acid secretion in unconscious rat and on basal acid secretion in conscious rat. *Journal of Medicinal Plants* **2009**;8:123-30.
 50. Eliassi A, Mandipour M, Kamalinejad M. Intra gastric effect of elaeagnus angustifolia L. fruit on gastric acid secretion in a rat pylorus - Ligated model. *Journal of Medicinal Plants* **2008**;7:82-91.
 51. Watkinson G, Hopkins A, Akbar F. The therapeutic efficacy of misoprostol in peptic ulcer disease. *Postgrad Med J* **1987**;64:60-77.
 52. Ghosh PK, Gaba A. Phyto-extracts in wound healing. *J Pharm Pharm Sci* **2013**;16:760-820.

53. Natanzi MM, Pasalar P, Kamalinejad M, Dehpour AR, Tavangar SM, Sharifi R, *et al*. Effect of aqueous extract of *Elaeagnus angustifolia* fruit on experimental cutaneous wound healing in rats. *Acta Med Iran* **2012**;50:589-96.
54. Guo Sa, DiPietro LA. Factors affecting wound healing. *J Dent Res* **2010**;89:219-29.
55. Burgess C. Topical vitamins. *J Drugs Dermatol* **2008**;7:s2-s6.
56. Biyik H. Antimicrobial activity of the ethanol extracts of some plants natural growing in Aydin, Turkey. *Afr J Microbiol Res* **2010**;4:2318-23.
57. Hambaba L, Boudjellal K, Abdeddaim M, Aberkane MC, Boudiaf K. In vitro investigation of antimicrobial and antioxidant activities of *Elaeagnus angustifolia* L. fruit's extracts. *Phytotherapie* **2012**;1-7. doi: 10.1007/s10298-012-0737-7.
58. Sharma N, Maiti S, Koley K. Studies on the incidence of sub clinical mastitis in buffaloes of Rajnandgaon district of Chhattisgarh state. *Vet Pract* **2004**;5:123-4.
59. Roberts D. Microbiological aspects of goat's milk. A Public Health Laboratory Service survey. *J Hyg (Lond)* **1985**;94:31-44.
60. Okmen G, Turkcan O. The antibacterial activity of *elaegnus angustifolia* l. Against mastitis pathogens and antioxidant capacity of the leaf methanolic extracts. *J Anim Vet Adv* **2013**;12:491-6. doi: 10.3923/javaa.2013.491.496.
61. Bucur L, Badea V, Istudor V. Antibacterial activity for two soft extracts of *Elaeagnus angustifolia* L. *Archives of the Balkan Medical Union* **2006**;41:127-31.
62. Khan SU, Khan AU, Shah AUHA, Shah SM, Hussain S, Ayaz M, *et al*. Heavy metals content, phytochemical composition, antimicrobial and insecticidal evaluation of *Elaeagnus angustifolia*. *Toxicol Ind Health* **2016**;32:154-61.
63. Okmen G, Turkcan O. A study on antimicrobial, antioxidant and antimutagenic activities of *Elaeagnus angustifolia* L. leaves. *Afr J Tradit Complement Altern Med* **2014**;11:116-20.
64. Bahraminejad S, Abbasi S, Amiri R. In Vitro Antifungal Activity of Various Methanolic Plant Extracts Against *Alternaria Solani*. *Iran J Pharm Res* **2013**;12:1470.
65. Malik F, Hussain S, Mirza T, Hameed A, Ahmad S, Riaz H, *et al*. Screening for antimicrobial activity of thirty-three medicinal plants used in the traditional system of medicine in Pakistan. *Journal of Medicinal Plants Research* **2011**;5:3052-60.
66. Wang Y, Guo T, Zhao C, Zhao P, Fan M. Changes in total phenolic and flavonoid contents and antioxidant activities of the fruit from *Elaeagnus angustifolia* during an 80-day study period. *Agro Food Ind Hi Tech* **2014**;25:7-10.
67. Shi C, Sun Z, Xie B. Extraction, purification and antioxidant of proanthocyanidins of *Elaeagnus angustifolia* L. Jujubes. *Nongye Gongcheng Xuebao/Transactions of the Chinese Society of Agricultural Engineering* **2006**;22:158-61.
68. Faramarz S, Dehghan G, Jahanban-Esfahlan A. Antioxidants in different parts of oleaster as a function of genotype. *Bioimpacts* **2015**;5:79-85.
69. Bucur L, Țarălungă G, Istudor V, Badea V. Pharmacological studies on *Elaeagnus angustifolia* L. tinctures and extracts. Note 2. Experimental research about the flowers tincture antitumoral activity. *Archives of the Balkan Medical Union* **2008**;43:107-12.
70. Giardi MT, Rea G, Berra B. *Bio-Farms for Nutraceuticals: Functional Food and Safety Control by Biosensors*. Springer Science & Business Media; **2011**.
71. Ou B, Hampsch-Woodill M, Prior RL. Development and validation of an improved oxygen radical absorbance capacity assay using fluorescein as the fluorescent probe. *J Agric Food Chem* **2001**;49:4619-26.
72. Yalcin G, Sogut O. Antioxidant capacity of *Elaeagnus angustifolia* L. and investigation of eosin y as the fluorescent probe in ORAC method. *J Food Agric Environ* **2014**;12:51-4.
73. Yalcin G, Sogut O. Influence of extraction solvent on antioxidant capacity value of oleaster measured by ORAC method. *Nat Prod Res* **2014**;28:1513-7.
74. Benvenuti S, Pellati F, Melegari Ma, Bertelli D. Polyphenols, anthocyanins, ascorbic acid, and radical scavenging activity of Rubus, Ribes, and Aronia. *J Food Sci* **2004**;69:FCT164-FCT9.
75. Ya W, Shang-Zhen Z, Chun-Meng Z, Tao G, Jian-Ping M, Ping Z, *et al*. Antioxidant and Antitumor Effect of Different Fractions of Ethyl Acetate Part from *Elaeagnus angustifolia* L. *Adv J Food Sci Technol* **2014**;6:707-10.
76. Si CL, Qin PP, Lu YY, Wu L, Wang HH, Hui LF, *et al*. GC-MS analysis of chemical composition and free radical scavenging activity of *elaegnus angustifolia* bark. Harbin; 2011. p. 854-8.
77. Chen Q, Chen J, Du H, Li Q, Chen J, Zhang G, *et al*. Structural characterization and antioxidant activities of polysaccharides extracted from the pulp of *Elaeagnus angustifolia* L. *Int J Mol Sci* **2014**;15:11446-55.
78. Caliskan E, Elmastas M, Gokce I. Evaluation of antioxidant properties of *elaegnus angustifolia* flowers. *Asian J Chem* **2010**;22:2840-8.
79. Bucur L, Negreanu-Pirjol T, Giurginca M, Istudor V. Some new *Elaeagnus angustifolia* L. Extracts and the pharmaceutical products' antioxidant activities determined by the chemiluminescence method. *Rev Roum Chim* **2008**;53:961-4.
80. Bendaikha S, Gadaut M, Harakat D, Magid A. Acylated flavonol glycosides from the flower of *Elaeagnus angustifolia* L. *Phytochemistry* **2014**;103:129-36.
81. Wang Y, Fan M, Li J, Guo T. Antitumor effect of edible part of *elaegnus angustifolia* L. in vivo and in vitro. *Journal of Chinese Institute of Food Science and Technology* **2013**;13:26-31.
82. Bucur L, Stanciu G, Istudor V. The GC-MS analysis of *Elaeagnus angustifolia* L. flowers essential oil. *Revista de Chimie* **2007**;58:1027-9.
83. Creamer D, Sullivan D, Bicknell R, Barker J. Angiogenesis in psoriasis. *Angiogenesis* **2002**;5:231-6.
84. Badrhadad A, Kh P, Mansouri K. In vitro anti-angiogenic activity fractions from hydroalcoholic extract of *Elaeagnus angustifolia* L. flower and *Nepeta crispa* L. arial part. *Journal of Medicinal Plants Research* **2012**;6:4633-9.
85. Srirama R, Ramesha B, Ravikanth G, Uma Shaanker R, Ganeshiah K. Are plants with anti-cancer activity resistant to crown gall?: a test of hypothesis. *Curr Sci* **2008**;95:1407-8.
86. Gill CI, Boyd A, McDermott E, McCann M, Servili M, Selvaggini R, *et al*. Potential anti-cancer effects of virgin olive oil phenol on colorectal carcinogenesis models in vitro. *Int J Cancer* **2005**;117:1-7.
87. Abu-Dahab R, Afifi F. Antiproliferative activity of selected medicinal plants of Jordan against a breast adenocarcinoma cell line (MCF7). *Sci Pharm* **2007**;75:121.
88. Hosseinzadeh H, Ramezani M, Namjo N. Muscle relaxant activity of *Elaeagnus angustifolia* L. fruit seeds in mice. *J Ethnopharmacol* **2003**;84:275-8. doi: 10.1016/s0378-8741(02)00331-8.
89. Wasowski C, Marder M. Flavonoids as GABAA receptor ligands: the whole story? *J Exp Pharmacol* **2012**;4:9.
90. Mohammed FI, Al-Essa MK, Shafagoj YA, Afifi FU. Investigation of the direct effects of the alcoholic extract of *Elaeagnus angustifolia* L. (*Elaeagnaceae*) on dispersed intestinal smooth muscle cells of guinea pig. *Sci Pharm* **2006**;74:21-30. doi: 10.3797/scipharm.2006.74.21.
91. Semenov B, Novikov K, Spitsin A, Azev V, Kachala V. Diastereotopic synthesis of 1- and 1, 1-substituted 4-phenyl-2, 3, 4, 9-tetrahydro-1H- β -carboline. *Chem Nat Compd* **2004**;40:585-90.
92. Wang B, Qu H, Ma J, Sun X, Wang D, Zheng Q, *et al*. Protective Effects of *Elaeagnus angustifolia* Leaf Extract against Myocardial Ischemia/Reperfusion Injury in Isolated Rat Heart. *J Chem* **2014**;2014:6. doi: 10.1155/2014/693573.
93. Tamtaji OR, Taghizadeh M, Takhtfiroozeh SM, Talaei SAR. The effect of *elaegnus angustifolia* water extract on Scopalamine-Induced memory impairment in rats. *Majallahi Ilmi Pizhuhishii Danishgahi Ulumi Pizishki Khadamatii Bihdashtii Darmanii Zanjan* **2014**;22:101-11.
94. Akbarzadeh M, Zeinalzadeh S, Zolghadri J, Mohagheghzadeh A, Faridi P, Sayadi M. Comparison of *Elaeagnus angustifolia* extract and sildenafil citrate on female orgasmic disorders: A randomized clinical trial. *J Reprod Infertil* **2014**;15:190-8.

95. Khodakarm-Tafti A, Mehrabani D, Homafar L, Farjanikish G. Healing effects of *Elaeagnus angustifolia* extract in experimentally induced ulcerative colitis in rats. *J Pharmacol Toxicol* **2015**;10:29-35.
96. Khafagy E-S, Morishita M, Onuki Y, Takayama K. Current challenges in non-invasive insulin delivery systems: a comparative review. *Adv Drug Deliv Rev* **2007**;59:1521-46.
97. Kaouah F, Boumaza S, Berrama T, Trari M, Bendjama Z. Preparation and characterization of activated carbon from wild olive cores (oleaster) by H₃PO₄ for the removal of Basic Red 46. *J Clean Prod* **2013**;54:296-306. doi: 10.1016/j.jclepro.2013.04.038.
98. Mohajjel Nayebi AR, Barzegar Jalali M, Pourmohammad S. Study of analgesic effect of naproxen solid dispersions in crosspovidone and *Elaeagnus Angustifolia* fruit powder by using formalin test. *Bioimpacts* **2009**;15:125-32.
99. Mohammadi G, Barzegar-Jalali M, Valizadeh H, Nazemiyeh H, Barzegar-Jalali A, Shadbad MRS, et al. Reciprocal powered time model for release kinetic analysis of ibuprofen solid dispersions in oleaster powder, microcrystalline cellulose and crospovidone. *J Pharm Pharm Sci* **2010**;13:152-61.
100. Bucur L, Tarălungă G, Alexandrescu R, Negreanu T, Istudor V. Evaluation of biological activity of a dermatological preparation with *elaeanus angustifolia* flowers soft extract. *Rev Med Chir Soc Med Nat Iasi* **2007**;112:1098-103.
101. Bucur L, Hirjau V, Istudor V. *Elaeagnus angustifolia* L. flowers soft extract valorification in a dermatological preparation. Note 1. Development of dermatological preparation and preliminary quality control. *Farmacia* **2008**;56:215.
102. Taheri JB, Anbari F, Maleki Z, Boostani S, Zarghi A, Pouralibaba F. Efficacy of *Elaeagnus angustifolia* topical gel in the treatment of symptomatic oral lichen planus. *J Dent Res Dent Clin Dent Prospects* **2010**;4:29-32.
103. Talaie-Khozani T, Vojdani Z, Dehghani F, Heidari E, Kharazinejad E, Panjehshahin MR. Toxic effects of *Elaeagnus angustifolia* fruit extract on chondrogenesis and osteogenesis in mouse limb buds. *Tokai J Exp Clin Med* **2011**;36:63-70.
104. Khan JA, Khan MS. Metal analysis, phytotoxic, insecticidal and cytotoxic activities of selected medicinal plants of Khyber Pakhtunkhwa. *Pak J Pharm Sci* **2012**;25:51-8.
105. Quiralte J, Florido F, Arias de Saavedra J, Gómez A, Saenz de San Pedro B, Gonzalez E, et al. Olive allergen-specific IgE responses in patients with *Olea europaea* pollinosis. *Allergy* **2002**;57:47-52.
106. Carnés Sánchez J, Iraola V, Sastre J, Florido F, Boluda L, Fernández-Caldas E. Allergenicity and immunochemical characterization of six varieties of *Olea europaea*. *Allergy* **2002**;57:313-8.
107. Sastre J, Lluch-Bernal M, Bustillo A, Carnes J, Maranon F, Casanovas M, et al. Allergenicity and cross-reactivity of Russian olive pollen (*Elaeagnus angustifolia*). *Allergy* **2004**;59:1181-6.
108. Songnuan W. Wind-pollination and the roles of pollen allergenic proteins. *Asian Pac J Allergy Immunol* **2013**;31:261.
109. Fonia A, White IR, White JML. Allergic contact dermatitis to *Elaeagnus* plant (Oleaster). *Contact Dermatitis* **2009**;60:178-9. doi: 10.1111/j.1600-0536.2008.01485.x.
110. Shah JFF, Harner MJ, Tibbets TM. *Elaeagnus angustifolia* elevates soil inorganic nitrogen pools in riparian ecosystems. *Ecosystems* **2010**;13:46-61. doi: 10.1007/s10021-009-9299-4
111. Çakmakçı S, Topdaş EF, Kalın P, Han H, Şekerci P, P. Köse L, et al. Antioxidant capacity and functionality of oleaster (*Elaeagnus angustifolia* L.) flour and crust in a new kind of fruity ice cream. *Int J Food Sci Technol* **2015**;50:472-81. doi: 10.1111/ijfs.12637.
112. Shore PA, Giachetti A. Reserpine: basic and clinical pharmacology. In: *Handbook of psychopharmacology*. Springer; **1978**. p. 197-219.
113. Daglia M. Polyphenols as antimicrobial agents. *Curr Opin Biotechnol* **2012**;23:174-81.
114. Newman DJ, Cragg GM. Natural Products as Sources of New Drugs over the Last 25 Years. *J Nat Prod* **2007**;70:461-77.