

Platelet-derived microparticles enhance Ara-C-induced cell death in acute lymphoblastic leukemia (Nalm-6)

Fariba Nikravesh^{1#}, Hossein Arezoomand^{1#}, Roohollah Mirzaee Khalilabadi¹, Maryam Nooshadokht², Hajar Mardani Valandani^{1*}

¹Department of Hematology and Laboratory Sciences, Faculty of Allied Medicine, Kerman University of Medical Sciences, Kerman, Iran

²Department of Medical Parasitology and Mycology, School of Medicine, Kerman University of Medical Sciences, Kerman, Iran

Article Info



Article Type:

Original Article

Article History:

Received: 15 Apr. 2024

Revised: 8 Jul. 2024

Accepted: 24 Jul. 2024

ePublished: 29 Oct. 2024

Keywords:

Acute lymphoblastic leukemia
Platelet-derived microparticle
Cytarabine
Leukemia microenvironment
Platelet


Abstract

Introduction: The current understanding highlights the intricate relationship between leukemic cells and their microenvironment, emphasizing the significant impact of environmental factors on chemotherapy resistance or sensitivity. Platelet-derived microparticles (PMPs) play a crucial role in facilitating intercellular communication, significantly contributing to the complex dynamics of cancer pathology and treatment outcomes. This study aims to investigate the cytotoxic and apoptotic effects of PMP, Ara-C, and their combinations on cancer cells, as well as their influence on the expression of critical genes like *Bax*, *Bcl-2*, *P21*, and *h-TERT* in the context of acute lymphoblastic leukemia (ALL) cell line (Nalm-6).

Methods: PMPs were isolated through centrifugation at varying speeds, and their concentration was determined using the BCA assay. The size and immunophenotypic characteristics of PMPs were analyzed using dynamic light scattering (DLS) and flow cytometry. The cytotoxic and apoptotic effects of PMP, Ara-C, and their combinations on Nalm-6 cells were assessed using the MTT assay, the trypan blue exclusion assay, and flow cytometry. Gene expression levels were analyzed using real-time PCR.

Results: According to our research findings, PMPs did not independently impact the viability and apoptosis of Nalm-6 cells; however, they synergistically augmented Ara-C's suppressive impact on viability and apoptosis. The MTT assay showed that both PMPs and Ara-C, whether administered alone or in combination, had a cytotoxic effect on the Nalm-6 cells. Furthermore, the combined treatment significantly affected the expression of *Bax*, *Bcl-2*, *P21*, and *h-TERT* genes.

Conclusion: Our study demonstrates that PMPs have the potential to improve the effectiveness of Ara-C chemotherapy in treating ALL. These findings contribute to a deeper understanding of the interplay between PMP and chemotherapy agents, offering potential insights for optimizing treatment strategies and improving patient outcomes in ALL.



Introduction

Acute lymphoblastic leukemia (ALL) is a fast-growing blood cancer caused by chromosome abnormalities and genetic changes that impact the growth and quick division of immature T or B cells.¹ These genetic changes lead to the accumulation of leukemic cells in the bone marrow and other tissues, suppressing normal hematopoiesis in the bone marrow.^{2,3} ALL accounts for 75% of leukemia cases in children and is the third most common type of

leukemia in adults. While ALL has a favorable prognosis and survival rates (around 80%) in children, it represents a severe condition in adults, with a survival rate below 40%.⁴ Chemotherapy in ALL consists of three phases: induction, consolidation, and maintenance.⁵ Ara-C is an effective chemotherapy drug for the treatment of ALL. This drug is often used in combination chemotherapy along with fluorouracil, methotrexate, and dexamethasone in solid tumors and leukemia.⁶ The impact of Ara-C stems from



*Corresponding author: Hajar Mardani Valandani, Email: hajarmardani61@gmail.com

These authors contributed equally in this project.



© 2025 The Author(s). This work is published by BioImpacts as an open access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/4.0/>). Non-commercial uses of the work are permitted, provided the original work is properly cited.

its active absorption into specific cells and subsequent conversion to its active form through metabolism. The rapid inactivation of Ara-C by plasma enzymes makes its plasma half-life about 10 minutes. As a result, for more effectiveness, Ara-C is prescribed in high doses.⁷ While low-dose Ara-C therapy is generally well tolerated, significant toxicities mainly occur with high-dose treatment. The dose-dependent toxicity is a notable limitation of Ara-C therapy.⁸ The objectives of ongoing ALL studies aim to optimize treatment strategies with less toxic interventions, enhance survival outcomes, and improve the overall quality of life for patients.⁹ Previous studies have demonstrated that, besides the dosage and combination of chemotherapy drugs, environmental factors significantly impact cancer cell biology and influence the overall effectiveness of cancer treatment.¹ Today, it is known that malignancies, including leukemia, are closely related to their environment, and platelets are an essential part of the cancer cell microenvironment.¹⁰ Studies demonstrate that cancer cells activate platelets, leading to the release of platelet granule contents and microparticles known as platelet-derived microparticles (PMPs).¹¹ PMPs are plasma membrane particles released from activated platelets during membrane budding. These particles constitute over 90% of microparticles found in the bloodstream.¹² PMPs, which carry various functional compounds of platelets, serve as potential mediators between platelets and cancer cells. PMPs are an especially interesting type of extracellular vesicles for evaluating cancer, as their increase has been observed in all cancer types, including stomach, lung, skin, CML, CLL, ALL, and others.¹³ PMPs affect growth and proliferation, apoptosis, and gene expression patterns in the target cells by transferring genetic and biochemical content.¹⁴⁻¹⁶ These findings show that PMPs can affect the biology of acute lymphoid leukemia cells and the effectiveness of chemotherapy drugs, improving risk-based treatments and reducing the dose and toxicity of drugs. Although various studies have investigated the dual effect of PMP in the progression and biology of cancer cells, the current knowledge in this field seems incomplete. The Previous study shows that Ara-C's impact on the expression of *Bax*, *Bcl-2*, *P21*, and *h-TERT* genes is crucial in its anti-cancer mechanism. It involves regulating the cell cycle, inducing apoptosis, and maintaining telomeres. Ara-C can boost the expression of proapoptotic genes like *Bax*, caspase-9, and caspase-3 at both mRNA and protein levels while suppressing the expression of the anti-apoptotic gene *Bcl-2* in renal carcinoma cells. Experiments have proven that Ara-C can impede the AKT/PI3K signaling pathway's activity, thereby promoting cell apoptosis.¹⁷⁻²⁰ In summary, while previous studies have investigated the role of PMPs in cancer biology and chemotherapy response, our study aims to provide novel insights into the specific mechanisms underlying the interaction between PMPs and cancer

cells, particularly in the context of Ara-C treatment for ALL. By examining the impact of PMPs, Ara-C, and their combination on critical parameters such as cell survival, apoptosis induction, and gene expression profiles such as *Bax*, *Bcl-2*, *P21*, and *h-TERT*. We hypothesize that PMPs enhance the apoptotic effects of Ara-C and modulate the expression of critical genes involved in cell survival and proliferation pathways. These findings may contribute to developing more effective and targeted therapies for ALL.

Materials and Methods

Preparation of platelet concentrates

Kerman Blood Transfusion Center provided expired platelet concentrate (PC) bags.

Separation of PMPs from PCs

A multi-step centrifugation technique was used to separate PMPs from PCs. To separate the cells (RBCs and WBCs), the PCs were transferred into 50 mL Falcon tubes and centrifuged at 2500 g for 10 minutes. This process was repeated to ensure the complete removal of any remaining cells. The resulting supernatant solution underwent centrifugation at 22000 g for 30 minutes to precipitate the PMP. The isolated PMP was then washed with PBS to eliminate any residual plasma and stored at -70 °C for future use.²¹

Characterization of isolated PMPs

Determination of the size of PMPs

Dynamic light scattering (DLS) is a scientific technique used to measure the dimensions and size distribution of particles or molecules in a liquid or solution. DLS relies on the concept that particles in a liquid undergo random, thermally induced movements known as Brownian motion. These movements cause fluctuations in the intensity of scattered laser light. During a DLS experiment, a sample containing particles is exposed to a focused laser beam, and a detector records the changes in scattered light intensity over time at a specific angle. The software integrated into the device is employed to determine the average size of the particles.

The Malvern Master Sizer 2000 laser diffraction apparatus, utilizing DLS technology, is utilized for sizing analysis. Within this setup, a laser beam with a 633 nm wavelength is aimed at the suspension housing microparticles. Subsequently, the scattered light is captured, and the instrument's software computes the dimensions of the microparticles.²²

Examining the surface marker of PMPs

Flow cytometry was used to validate the separation technique and confirm the cellular origin of the isolated PMPs by identifying platelet-specific markers CD42b and CD61 on the surface of the PMPs. 250 µL of PMP suspension was mixed with an equal volume of PBS (1:1 ratio in PBS). Subsequently, 100 µL of the diluted PMP

solution was combined with two test tubes and two control tubes, each containing 10 μ L of CD42b and CD61 antibodies, as well as 10 μ L of IgG-FITC/PE antibody targeting mouse cells. After a thirty-minute incubation period at room temperature in the dark, the samples were analyzed using a flow cytometry instrument from BD Bioscience, with data management carried out using FLOMAX software.²³

Determining the concentration of PMPs

The BCA method, which has been endorsed by the International Society of Extracellular Vesicles (ISEV), is a reliable technique for quantifying protein concentrations in various samples, including PMPs.²⁴ The approach employed in this technique is based on the ability of proteins to convert copper ions into a colored chelate complex. This complex can then be measured using spectrophotometry to determine its quantity. The total protein content of PMPs was assessed using the Micro BCA Protein Assay Kit (Pierce, Thermo Scientific, Rockford, IL, USA) by the manufacturer's guidelines. Bovine serum albumin (BSA) served as the standard for comparative purposes.²¹

Cell culture

The Nalm-6 cell line (Nalm-6, NCBI C212) was acquired from the Pasteur Institute collection located in Tehran, Iran. The Nalm-6 cells were cultured in RPMI-1640 medium, which was supplemented with 10% fetal bovine serum (FBS) (Gibco™ A3160402) and 1% antibiotic (Penicillin-Streptomycin Solution 100X, Biowest, L0022). Subsequently, the cells were incubated in a humidified incubator at a temperature of 37 °C with a 5% CO₂ atmosphere.²⁵

Calculation of the combination index and the dose reduction index

CI (combination index) between PMP and Ara-C was determined using CompuSyn Software (ComboSyn, Inc., Paramus, NJ, USA). CI to quantitatively depict synergism ($CI < 1$), additive effect ($CI = 1$), and antagonism ($CI > 1$).

The dose reduction index (DRI) quantifies how much the dosage of a combination can be decreased to attain a particular level of effect compared to the concentration of the individual drug used alone.²⁶

Trypan Blue assay

Nalm-6 cells were cultured in a 6-well plate at a density of 5×10^5 cells per well. The cells were then treated with different concentrations of PMP (200, 400, and 800 μ g/mL) and Ara-C (0.5, 1, 1.5, and 2 μ M) for 24, 48, and 72 hours to evaluate the apoptotic effect of these compounds on cell viability. The cell suspension underwent centrifugation, followed by placing the pellet in a serum-free complete medium. Subsequently, a mixture consisting of one part 0.4% trypan blue (Gibco™ 15250061) and one

part cell suspension was incubated for 2 minutes at room temperature. The total number of unstained (viable) and stained (non-viable) cells was manually counted using a Neuburger chamber and a light microscope (ECLIPSE E100, Nikon). Finally, the percentage of viable cells was determined by calculating "Viability (%) = viable cells / (viable cells + death cells) $\times 100$ ".²⁷

MTT assay

The MTT colorimetric assay was utilized to assess the cytotoxicity of PMP, Ara-C, and their combinations against cancer cell lines in vitro. This assay measures the capacity of viable cells to convert yellow tetrazolium salts into blue-colored formazan crystals. Nalm-6 cells were seeded at a density of 1×10^4 cells per well in 96-well plates and exposed to varying concentrations of PMP (200, 400, and 800 μ g/mL) and Ara-C (1 and 1.5 μ M) for durations of 24, 48, and 72 hours. Furthermore, viability evaluations were carried out for combinations of PMP/Ara-C, such as PMP (200 μ g/mL) + Ara-C (1 μ M), PMP (200 μ g/mL) + Ara-C (1.5 μ M), PMP (400 μ g/mL) + Ara-C (1 μ M), and PMP (400 μ g/mL) + Ara-C (1.5 μ M). The plate was then centrifuged at 700g for 10 minutes, and the supernatant was removed. The cells were then exposed to 100 μ L MTT solution (0.5 mg/mL; (M5655, Sigma) at 37 °C. After 4 hours, the colored formazan was dissolved in each well with 150 μ L DMSO, and the optical absorbance at 570 nm was gauged using an enzyme-linked immunosorbent assay (ELISA). The metabolic activity of the treated cells was quantified as a ratio of untreated cells utilized as a negative control.²⁸

Flow cytometry

The impact of PMP, Ara-C, and their respective combinations on initiating early and late apoptosis was assessed through annexin-V and propidium iodide (PI) staining. A total of 4×10^5 cells were seeded per well on a six-well cell culture plate and subjected to treatment with PMP (400 μ g/mL), Ara-C (1.5 μ M), or a combination of both (400 μ g/mL PMP and 1.5 μ M Ara-C). Subsequently, flow cytometry analysis was performed using the Annexin-V Apoptosis Detection Kit (Mab Tag, AnxF100PI), and the resulting data were analyzed using FlowJo version 7.6.1 software.²⁹

Gene expression analysis using real-time PCR

Real-time PCR was used to monitor changes in mRNA expression of *Bax*, *Bcl-2*, *P21*, and *h-TERT* genes. Total RNA was isolated using YTzol Pure RNA (Yekta Tajhiz Azma, YT9066) from untreated (control) and treated cells with 400 μ g/mL PMP, 1.5 μ M Ara-C, and PMP/Ara-C combination (400 μ g/mL PMP and 1.5 μ M Ara-C) following a 48-hour treatment period. The purity and concentration of the extracted RNA were determined by analyzing the ratio of light absorption at wavelengths of 260 nm and

280 nm using a NanoDrop 1000 Spectrophotometer. Moreover, the quality of the extracted RNA was assessed by subjecting RNA molecules to electrophoresis on an agarose gel for separation. Afterward, cDNA was synthesized according to the manufacturer's instructions (Thermo Scientific Fermentas, K1622). As described previously, real-time PCR was used to monitor changes in the mRNA expression of the desired genes.²⁸ Table 1 shows the nucleotide sequences of the primers used in real-time RT-PCR.

Statistical analysis

The statistical analysis for this research was conducted using SPSS26 software to assess the assumptions, while GraphPad Prism 8 software was utilized for generating graphs. A significance level of 5% (alpha error) was set as the threshold for either rejecting or confirming the null hypothesis. All mean comparisons were conducted using two-way tests. Statistical significance was determined with a *P* value below 0.05. For the analysis of cell survival, metabolic activity, apoptosis, and gene expression data, two-way repeated measures ANOVA was employed in this study. Sample size calculation was performed using G-Power software, considering an alpha error of 5%, a

test power of 60%, an effect size of 50%, and accounting for four treatments measured at two different time points. The calculated sample size for each group was 3.

Results

Determination of the size of PMPs

The average size of the isolated PMPs was 200 nm for 90%, as shown in Fig. 1.

Phenotypical characterization of PMPs

The immunophenotypic characterization revealed that 97% of the PMPs expressed CD61 (Fig. 2a), and 95% expressed CD42b (Fig. 2b) on their surface.

The impact of Ara-C on the cell mortality of Nalm-6 cells was increased after co-incubation with PMP

The effects of different concentrations of Ara-C (0.5, 1, 1.5, and 2 μ M) and PMP (200, 400, and 800 μ g/mL), both individually and in combination, on cell viability were studied at 24, 48, and 72 hours. According to Fig. 3a, PMP treatment did not affect the viability of Nalm-6 cells. In contrast, as illustrated in Fig. 3b, the Nalm-6 cell viability decreased following exposure to Ara-C. Notably, co-incubation of cells with the combination of Ara-C and PMP for 48 h of treatment reduced cell viability more effectively than to Ara-C alone, as illustrated in Fig. 3c.

The cytotoxic impact of Ara-C and PMP on Nalm-6 cells intensified following concurrent incubation

The cytotoxic impact of PMP and Ara-C at different concentrations (1, 1.5 μ M for Ara-C; 200, 400, and 800 μ g/mL for PMP) and their combination was assessed. PMP and Ara-C demonstrated independent cytotoxic effects on Nalm-6 treated cells (Fig. 4a, b). The concurrent treatment of PMP and Ara-C synergistically amplified the cytotoxic effects on the Nalm-6 cells (Fig. 4c). Synergistic effects ($CI < 1$) were observed for two combination doses: PMP (400 μ g/mL) + Ara-C (1 μ M) and PMP (400 μ g/mL) + Ara-C (1.5 μ M). An additive effect ($CI = 1$) was

Table 1. PCR primer sequences

Gene	Sequence 5'-3'	T _m	nt #
GAPDH-FW	CCAACCGCGAGAAGATGA	50.3	18
GAPDH-Rv	TCCATCACGATGCCAGTG	50.3	18
Bax-FW	AGGATCGAGCAGGGCGAATG	62.5	20
Bax-Rv	TCAGCTTCTTGGTGGACGCA	60.5	20
P21-FW	CCTGTCACTGTCTGTACCT	54.4	21
P21-Rv	GCGTTTGGAGTGGTAGAAATCT	53	22
Bcl-2-FW	ATCGCCCTGTGGATGACTGAG	55.9	21
Bcl-2-Rv	CAGCCAGGAGAAATCAAACAGAG	55.9	23
h-TERT -FW	CGGAAGAGTGCTGGAGCAA	60	20
h-TERT -Rv	GGATGAAGCGGAGTCTGGA	59.3	19




Fig. 1. Determining the average size of microparticles derived from platelets by Malvern Master Sizer 2000. The size dispersion of the separated PMP was evaluated within a span of 200 nm.




Fig. 2. Flow cytometry analysis revealed a high expression of CD61 (a) molecules (97.6%) and CD42b (b) molecules (95%) on the PMP.




Fig. 3. The viability of Nalm-6 cells was assessed for 24, 48, and 72 h to determine the impact of different doses of (a) PMP, (b) Ara-C, and (c) the combination of PMP and Ara-C. The assessment was conducted using trypan blue staining, and the results were presented as mean \pm standard error (SE) with a sample size of 3. Interestingly, the various concentrations of PMP (200, 400, and 800 μ g/mL) did not show any significant effect on Nalm-6 cell viability. However, the effects of Ara-C (0.5, 1, 1.5, and 2 μ M) were found to be dose- and time-dependent. This suggests that the viability of Nalm-6 cells was influenced by the concentration of Ara-C and the duration of exposure. Furthermore, the impact of PMP and Ara-C in combination on Nalm-6 cell viability was evaluated after 48 h. The results indicated that the combination dose significantly reduced cell viability compared to Ara-C alone. This suggests a potential synergistic effect between PMP and Ara-C in inhibiting the viability of Nalm-6 cells (** $P < 0.01$, *** $P < 0.001$ relative to untreated cells).




Fig. 4. The cytotoxic effect of PMP (a) and Ara-C (b) on Nalm-6 cells was conducted at 24, 48, and 72 hours following treatment. Both PMP and Ara-C demonstrated toxicity effects. Additionally, (c) the collective impact of PMP and Ara-C was evaluated in comparison to separate dosages following a 48-hour period. PMP and Ara-C exhibited distinct cytotoxic impacts on Nalm-6 cells. The concurrent administration of PMP and Ara-C demonstrated a significant influence. Significant effects of the drug combination were noted in four specific doses (PMP and Ara-C: 200 + 1, 200 + 1.5, 400 + 1, 400 + 1.5) (* $P < 0.05$, ** $P < 0.01$, *** $P < 0.001$, relative to untreated cells).

noted for PMP (200 μ g/mL) + Ara-C (1 μ M), and an antagonistic effect ($CI > 1$) for PMP (200 μ g/mL) + Ara-C (1.5 μ M) (Fig. 5). Table 2 displays the CI and DRI values for the combination treatments, with the combination of 1.5 μ M Ara-C and 400 μ g/mL PMP showing strong synergistic effects, chosen for subsequent experiments.

PMP enhanced Ara-C apoptosis induced by Ara-C

The impact of Ara-C alone and in combination with PMP on programmed cell death was explored. Fig. 6a shows the control group that was not exposed to any treatment. Fig. 6b illustrates that the co-incubation of Nalm-6 cells with PMP for 48 hours did not lead to a notable effect on




Fig. 5. The synergistic effects of the combined therapy involving PMP and Ara-C on Nalm-6 cells were evaluated through the analysis of the CI versus FA Curve. Different combinations of PMP and Ara-C were administered, and their effect on the metabolic activity of the cells was evaluated using the MTT assay to generate CI vs FA data. (a) CI values below 1 indicate a synergistic effect, values at 1 represent an additive effect (shown by the solid line), and values above 1 indicate antagonism. It is worth noting that the combination doses of PMP (400 µg/mL) + Ara-C (1 µM) and PMP (400 µg/mL) + Ara-C (1.5 µM) exhibited significant synergistic effects. (b) Furthermore, the dose-normalized isobologram analysis demonstrated that points below the effect line indicate synergism, revealing a synergistic effect for two combination points in Nalm-6 cells.




Fig 6. Nalm-6 cells were subjected to various treatments, including control (untreated) (a), PMP: 400 µg/mL (b), Ara-C: 1.5 µM (c), and PMP: 400/Ara-C: 1.5 (d) for a duration of 48 h. Flow cytometry was utilized to evaluate Annexin-V and Annexin-V plus Propidium Iodide (PI) expression. Following a 48 h co-incubation with PMP, there was no significant impact on cell apoptosis in Nalm-6 cells. Conversely, Annexin V +/PI+ cells notably increased in the Ara-C treated group. It is apparent that PMP significantly enhanced the apoptotic effect of Ara-C on Nalm-6 cells (* $P < 0.05$, ** $P < 0.01$, *** $P < 0.001$, **** $P < 0.0001$ relative to untreated cells).

Table 2. CI and DRI for drug combination by Ara-C and PMP

Ara-C (µM)	DRI	PMP (µg/mL)	DRI	CI value (At inhibition of 50%)
1.0	1.5	200	2.757	1.029
1.0	2.231	400	1.889	0.977
1.5	1.172	200	3.129	1.172
1.5	3.32	400	3.568	0.581

cell apoptosis. Annexin V + /PI+ cells in the Ara-C treated group increased significantly (Fig. 6c). According to Fig. 6d, 48 h co-treatment of Nalm-6 cells with the combination

of 1.5µM Ara-C plus 400µg/ml PMP induced cell death more than the Ara-C group. These findings suggest that PMP enhances the apoptotic effect of Ara-C.

The impact of the PMP/Ara-C combination on gene expression

The expression of genes associated with cell death was examined further to study the effect of the PMP on Ara-C-dependent apoptosis. Ara-C upregulated the expression of the *Bax* gene, as illustrated in Fig. 7a, while PMP had no impact on *Bax* gene expression. The combination of




Fig 7. Fold change gene expression. **(a)** The graph presents that PMP does not affect *Bax* gene expression, but Ara-C upregulates the *Bax* gene (proapoptotic). **(b)** Also, PMP does not affect *Bcl-2* gene expression, but Ara-C downregulates the *Bcl-2* gene (anti-apoptotic). After 48 h of treatment, a combination of PMP and Ara-C induces more significant changes in gene expression compared to single doses in Nalm-6 cells. **(c)** *P21* gene expression increases in all three groups, although it is more significant in the combined group than in the other two. **(d)** Also, PMP treatment increased *h-TERT* gene expression in Nalm-6 cells, but Ara-C treatment decreased *h-TERT* gene expression. Furthermore, the combined dosage resulted in a substantial reduction in *h-TERT* gene expression (* $P < 0.05$, ** $P < 0.01$, *** $P < 0.001$, relative to untreated cells).

PMP and Ara-C resulted in a higher *Bax* gene expression compared to Ara-C treatment alone, as indicated by the results. The results (Fig. 7b) revealed that PMP did not affect the *Bcl-2* gene in Nalm-6 cells. However, Ara-C decreased its expression. When cells were co-incubated with PMP and Ara-C, the expression of this gene was less than in the control group, Ara-C alone, and PMP alone. These results showed that PMP increases Ara-C-induced apoptosis in Nalm-6 cells. To further understand the effect of the PMP/Ara-C combination on the molecular mechanism of cell population reduction, the expression of the proliferative gene (*P21*) and the gene associated with telomerase (*h-TERT*) was studied. As seen in Fig. 7c, *P21* gene expression increases in all three groups, although it is more significant in the combined group than in the other two. PMP treatment increased *h-TERT* gene expression in Nalm-6 cells, but Ara-C treatment decreased *h-TERT*

gene expression. Furthermore, co-treatment of cells with PMP plus Ara-C resulted in a substantial reduction in *h-TERT* gene expression (Fig. 7d).

Discussion

Treatment resistance poses a significant challenge for individuals with ALL despite advancements in pharmacological therapies.³⁰ While leukemia treatments have become more effective and less toxic over time, mortality rates persist as a concern. Combined therapy is often recommended to reduce cytotoxic dosages while enhancing treatment efficacy. Ara-C has demonstrated effectiveness in treating various hematologic cancers, including ALL.³¹ Numerous studies have explored the potential of combination therapy to improve ALL treatment, involving Ara-C and other drugs.³² Various factors within the tumor microenvironment play crucial

roles in determining the sensitivity of cells to therapeutic drugs in different types of cancers.³³ Platelets contain growth factors, inflammatory cytokines, and immune factors that modify tumor biology.³⁴ The intricate roles of platelet cargo in cancer are still not fully understood, with their influence probably varying based on factors like tumor type and stage.³⁵ PMPs have attracted considerable interest due to their potential involvement in facilitating communication between platelets and cancer cells.³⁶ PMPs influence cancer cell behavior by delivering a varied payload of nucleic acids, proteins, and lipids.³⁷⁻³⁹ There has been a significant increase in research focusing on PMP and its roles in various cellular functions and disease states.⁴⁰ Recently, considerable investigation has focused on the impact of PMPs in combination with chemotherapy drugs, mainly through in vitro studies. For example, Cacic et al demonstrated that PMPs protect the THP-1 cell line against daunorubicin-induced apoptosis.⁴¹ Additionally, in 2023, Gharib et al studied the resistance of cell-related chronic lymphocytic leukemia cells treated with PMP against cytarabine.⁴² These findings collectively advance our understanding of the intricate interplay between PMPs and chemotherapy agents in cancer treatment. Despite numerous studies on the interaction of PMPs and cancer, our understanding of their effectiveness in combination with chemotherapy drugs remains largely inadequate and necessitates further comprehensive research. In this research, we explore the cytotoxic and apoptotic effects of PMP, Ara-C, and their combinations on Nalm-6 cells, as well as their influence on the expression of critical genes like *Bax*, *Bcl-2*, *P21*, and *h-TERT*. Our study confirms the tumor-inhibiting effect of PMPs on cell lines in vitro. The cytotoxic effect of PMP on Nalm-6 cells and reduction in metabolic activity of Nalm-6 cells after PMP treatment is consistent with previous research indicating a potential anti-leukemic influence associated with PMPs. Vismora et al showed that PMPs significantly reduced the metabolic activity of breast cancer cells without notably affecting their viability.⁴⁴ Additionally have been shown that PMPs inhibit mitochondrial function in the lung and colon cells in mice.⁴⁵ Our study demonstrates that both PMP and Ara-C exhibit cytotoxicity and reduce metabolic activity in treated Nalm-6 cells. Moreover, the concurrent treatment with PMP and Ara-C synergistically enhances the cytotoxic effects on cells. Studies have shown varying results regarding the impact of PMPs on cancer metabolism. A study examined the influence of PMPs on cell lines associated with chronic lymphocytic leukemia, revealing a significant metabolic change from early to later stages.⁴² Furthermore, the CompuSyn analysis demonstrated significant synergistic effects on Nalm-6 cells with two combination doses of Ara-C and PMP (400 µg/mL + 1 µM and 400 µg/mL + 1.5 µM). These findings highlight the potential synergistic impact of PMP and Ara-C combination therapy on Nalm-6 cells, suggesting

a promising avenue for further investigation. CD36, GNAI2, GNB2, PPP2R1A, PRKCA, PRKACB, PPP2CA, PRKAR1A, ITPR1, FASN, GNAQ, GNG5, and TKT are among the proteins associated with PMPs that play crucial roles in diverse signaling pathways and regulate the metabolic activity of target cells.⁴⁶ Additionally, another study suggests that PMPs may influence metabolic activity by transferring genetic materials, such as miRNAs.⁴⁵

Co-incubation of Nalm-6 cells with PMPs increased their sensitivity to Ara-C at a concentration of 1.5 µM Ara-C plus 400 µg/ml PMP. Studies have shown PMPs potentially influence cellular signaling pathways that sensitize cells to apoptosis. Baj-Krzyworzeka and colleagues conducted a study examining the effects of PMPs on hematological cell lines, revealing that PMPs activate MAPKp42/44 and PI-3K-AKT in hematopoietic cells.⁴⁷ Moreover, Fateme Yari et al. showed that PMP induces apoptosis in PBMCs isolated from pre-B acute lymphoblastic leukemia.⁴³ Several studies have observed that PMPs affect cancer cell apoptosis induced by chemotherapy drugs. For instance, Cacic and colleagues demonstrated that co-incubation of THP-1 cells with PMPs elevated levels of miR-125a, miR-125b, and miR-199, which protected THP-1 cells against daunorubicin-induced cell death.⁴⁸ Recent findings also highlight the role of PMPs in the metabolic reprogramming of CLL cells, contributing to their increased resistance to chemotherapy drugs such as cytarabine, venetoclax, and plumbagin.⁴²

To further investigate the underlying mechanisms, we assessed the impact of PMPs, Ara-C, and their combination on the expression of apoptotic genes like *Bax* and *Bcl-2*. *Bcl-2* and *Bcl-XL*, members of the anti-apoptotic *Bcl-2* family, impede cell death by inhibiting caspase activation crucial for apoptosis or by preventing the release of apoptotic factors like cytochrome C and AIF from the mitochondria.⁴⁹ In this study, the expression of the *Bcl-2* gene in cells treated with PMPs did not show significant differences compared to the control group. However, gene expression decreased in cells exposed to Ara-C alone and in combination with PMPs. Importantly, PMPs did not affect the expression of the *Bax* gene. Nevertheless, *Bax* gene expression was higher in cells treated with both PMPs and Ara-C compared to cells treated with Ara-C alone. The analysis of *Bax* and *Bcl-2* gene expressions provides a plausible explanation for the increased susceptibility of Nalm-6 cells to chemotherapy drugs following treatment with PMPs and Ara-C.

Previous research has shown that PMPs transfer their genetic material, including miRNAs, to target cells, influencing gene expression and the apoptotic response in laboratory settings.⁵⁰ PMPs contain a rich reservoir of proteins, lipids, second messengers, and various RNA species. These components are transferred to target cells, altering cell signaling pathways and modulating

the cellular response to drugs.⁵¹ However, the exact mechanisms underlying these changes in drug response are not fully understood and require further investigation.

We also examined the effects of PMP, Ara-C, and the combination of these two on the expression of *P21* and *h-TERT* genes. The study findings revealed that PMP treatment led to an increase in the expression of the *P21* gene, with molecular data consistently aligning with the observed decrease in cell proliferation in cell culture. The results revealed that treating Nalm-6 cells with Ara-C, either alone or in combination with PMPs, enhanced the expression of the *P21* gene. The influence of PMPs on proliferation control and cell population was assessed using Neobar slides and a cell counter. The results showed a significant decrease in cell population after treatment with both PMPs and Ara-C. The inhibitory effect of Ara-C was further enhanced when combined with PMP treatment.

Several studies have highlighted the potential tumor-suppressing properties of PMPs, demonstrating their ability to inhibit tumor progression in mouse models of lung and colon cancer.⁴⁵ Furthermore, in laboratory experiments, PMPs have been observed to suppress the growth of cancer cell lines, such as THP-1 and HUVEC.^{37,41} Contrary to the findings in the study conducted by Mirzaee et al, it was demonstrated that PMPs reduce the expression of *P21* and *P53* genes in Mesenchymal Stem Cells, leading to increased cell proliferation. The differing effects observed in different cell lines suggest that the impact of PMPs may vary depending on the specific type of target cell.²³

The *P21* gene inhibits cancer cell proliferation through pathways that involve both *P53*-dependent and independent mechanisms. These pathways include regulating metabolic pathways and triggering cellular senescence.⁵² Ensuring the longevity of telomeres through the activation of telomerase, particularly *h-TERT*, is essential for the continuous renewal of cells, which is a hallmark of cancer. The regulation of *h-TERT* transcription is critical for activating telomerase in human cancers. This underscores the importance of mechanisms that govern telomere maintenance in cancer development and their potential as indicators in oncology.⁵³ While telomerase activity is typically observed in germline, hematopoietic, and rapidly renewing cells, this study sheds light on the intriguing interplay between PMP and *h-TERT* gene expression, demonstrating that PMP can provoke a notable increase in *h-TERT* gene expression levels. In the investigation by Samareh et al, real-time PCR analysis revealed a remarkable over threefold surge in *h-TERT* gene expression upon treatment with PMP, compared to the control group.⁵³ Ara-C appears to cause a reduction in the expression of this gene. Interestingly, when Ara-C is given with PMP, it lessens the inhibitory effect of Ara-C on *h-TERT* gene expression. However, the

precise mechanism behind this observed phenomenon is still unclear and requires additional investigation. Our study underscores the influence of PMPs on cancer biology. PMPs reduce cell proliferation and have a cytotoxic effect on cells, impacting gene expression both alone and in on with Ara-C. These results underscore the complex relationship between PMPs and cancer cell behavior, providing valuable insights for enhancing therapeutic strategies in cancer treatment.

Furthermore, our study is consistent with previous investigations,⁵⁴ contrasting with reports of PMPs enhancing cancer progression in other studies.^{15,47} This dual potential of PMPs to either promote cancer progression or suppress tumor growth suggests a complex interaction between cancer cells and PMPs, likely influenced by factors such as the conditions of PMP creation, cell type, and cancer stage.

It is important to clarify that this study specifically investigated the interaction between PMPs and Nalm-6 cells, as well as the combined effect of PMPs and Ara-C, in an in vitro setting.

It is worth noting that results obtained in vivo may vary due to the influence of environmental factors and the interplay between the cell line and the immune system. Moreover, the specific contents of PMPs and their mechanism of action remain crucial considerations for understanding their effects. However, these aspects were not investigated in our study, which aimed to explore the overall impact of PMPs.

To strengthen the validity of our findings, future experiments should confirm our results using a wider variety of leukemia cell lines and additional chemotherapy drugs. This broader approach will help ensure the reliability and applicability of our conclusions across different contexts and treatments in leukemia research. Additionally, including samples from patients in the early stages of leukemia would provide valuable insights into the potential applicability of PMPs in diverse clinical contexts. This comprehensive approach is essential for gaining a deeper understanding of the therapeutic potential of PMPs in cancer treatment.

Advancing our understanding of the dynamic relationship between cancer cells and PMPs holds significant implications for potential therapeutic interventions. However, additional research is needed to uncover the precise mechanisms underlying PMPs' role in modulating cancer cell behavior. Specifically, efforts should focus on elucidating the specific molecules responsible for these effects.

Conclusion

Our study offers a new point of perspective on the interaction between PMPs and cancer, indicating that PMPs have the potential to impact cancer cell behavior and trigger unique reactions. These findings reinforce

Research Highlights

What is the current knowledge?

- PMPs are known to affect cancer cell biology and chemotherapy response.
- Ara-C is a standard chemotherapy drug for ALL with dose-dependent toxicity.

What is new here?

- PMPs enhance Ara-C-induced apoptosis in Nalm-6 cells.
- PMPs modulate the expression of critical genes Bax, Bcl-2, P21, and h-TERT.

the expanding evidence on the anti-tumor properties of PMPs and emphasize the necessity for comprehensive investigations to grasp their underlying mechanisms fully. Our research underscores the significance of examining the influence of environmental factors like PMPs in cancer biology, which could ultimately pave the way for developing more efficacious treatments for leukemia and various other cancer types.

Acknowledgments

We sincerely thank the Kerman Blood Transfusion Center.

Authors' Contribution

Conceptualization: Hajar Mardani Valandani.

Formal analysis: Fariba Nikravesh.

Funding acquisition: Hajar Mardani Valandani.

Investigation: Fariba Nikravesh, Hossein Arezoomand, Maryam Nooshadokht.

Methodology: Hajar Mardani Valandani, Fariba Nikravesh, Hossein Arezoomand.

Project administration: Hajar Mardani Valandani, Fariba Nikravesh.

Supervision: Hajar Mardani Valandani.

Writing—original draft: Fariba Nikravesh, Hossein Arezoomand.

Writing—review & editing: Hajar Mardani Valandani, Fariba Nikravesh, Hossein Arezoomand, Roohollah Mirzaee Khalilabadi.

Competing Interests

The authors declare that they have no conflict of interest.

Ethical Statement

This study was performed in line with the principles of the Declaration of Helsinki. Approval was granted by the Ethics Committee of the Vice Chancellor for Research and Technology of Kerman University of Medical Sciences (IR.KMU.REC.1399.420). Informed consent was obtained from all individual participants included in the study.

Funding

This study was supported by the Vice Chancellor for Research and Technology of Kerman University of Medical Sciences (Grant number 400001101). The authors have no relevant financial or non-financial interests to disclose.

References

1. Cao Y, Wu C, Song Y, Lin Z, Kang Y, Lu P, et al. Cyr61 decreases cytarabine chemosensitivity in acute lymphoblastic leukemia cells via NF- κ B pathway activation. *Int J Mol Med* **2019**; 43: 1011-20. doi: 10.3892/ijmm.2018.4018.
2. Jabbour E, O'Brien S, Konopleva M, Kantarjian H. New insights into the pathophysiology and therapy of adult acute lymphoblastic leukemia. *Cancer* **2015**; 121: 2517-28. doi: 10.1002/cncr.29383.
3. Mei L, Ontiveros EP, Griffiths EA, Thompson JE, Wang ES, Wetzler M. Pharmacogenetics predictive of response and toxicity in acute lymphoblastic leukemia therapy. *Blood Rev* **2015**; 29: 243-9. doi: 10.1016/j.blre.2015.01.001.
4. Inaba H, Mullighan CG. Pediatric acute lymphoblastic leukemia. *Haematologica* **2020**; 105: 2524-39. doi: 10.3324/haematol.2020.247031.
5. Terwilliger T, Abdul-Hay M. Acute lymphoblastic leukemia: a comprehensive review and 2017 update. *Blood Cancer J* **2017**; 7: e577. doi: 10.1038/bcj.2017.53.
6. Li Z, Guo JR, Chen QQ, Wang CY, Zhang WJ, Yao MC, et al. Exploring the antitumor mechanism of high-dose cytarabine through the metabolic perturbations of ribonucleotide and deoxyribonucleotide in human promyelocytic leukemia HL-60 cells. *Molecules* **2017**; 22: 499. doi: 10.3390/molecules22030499.
7. Reese ND, Schiller GJ. High-dose cytarabine (HD araC) in the treatment of leukemias: a review. *Curr Hematol Malig Rep* **2013**; 8: 141-8. doi: 10.1007/s11899-013-0156-3.
8. Di Francia R, Crisci S, De Monaco A, Cafiero C, Re A, Iaccarino G, et al. Response and toxicity to cytarabine therapy in leukemia and lymphoma: from dose puzzle to pharmacogenomic biomarkers. *Cancers (Basel)* **2021**; 13: 966. doi: 10.3390/cancers13050966.
9. Lato MW, Przysucha A, Grosman S, Zawitkowska J, Lejman M. The new therapeutic strategies in pediatric T-cell acute lymphoblastic leukemia. *Int J Mol Sci* **2021**; 22: 4502. doi: 10.3390/ijms22094502.
10. Repsold L, Pool R, Karodia M, Tintinger G, Joubert AM. An overview of the role of platelets in angiogenesis, apoptosis and autophagy in chronic myeloid leukaemia. *Cancer Cell Int* **2017**; 17: 89. doi: 10.1186/s12935-017-0460-4.
11. Lazar S, Goldfinger LE. Platelet microparticles and miRNA transfer in cancer progression: many targets, modes of action, and effects across cancer stages. *Front Cardiovasc Med* **2018**; 5: 13. doi: 10.3389/fcvm.2018.00013.
12. Boilard E. Platelet-derived nanomedicine targets cancer. *Blood* **2017**; 130: 561-2. doi: 10.1182/blood-2017-05-786327.
13. Saber SH, Ali HE, Gaballa R, Gaballah M, Ali HI, Zerfaoui M, et al. Exosomes are the driving force in preparing the soil for the metastatic seeds: lessons from the prostate cancer. *Cells* **2020**; 9: 564. doi: 10.3390/cells9030564.
14. Evangelista FC, Ferrão AL, Duarte RC, Gomes LC, Alves LC, Campos FM, et al. Circulating microparticles and thrombin generation in patients with chronic lymphocytic leukemia. *Braz J Pharm Sci* **2022**; 58: e19407. doi: 10.1590/s2175-97902022e19407.
15. Liang H, Yan X, Pan Y, Wang Y, Wang N, Li L, et al. MicroRNA-223 delivered by platelet-derived microvesicles promotes lung cancer cell invasion via targeting tumor suppressor EPB41L3. *Mol Cancer* **2015**; 14: 58. doi: 10.1186/s12943-015-0327-z.
16. Noulisri E. Effects of cell-derived microparticles on immune cells and potential implications in clinical medicine. *Lab Med* **2021**; 52: 122-35. doi: 10.1093/labmed/lmaa043.
17. Anderson NM, Harrold I, Mansour MR, Sanda T, McKeown M, Nagykarly N, et al. BCL2-specific inhibitor ABT-199 synergizes strongly with cytarabine against the early immature LOUCY cell line but not more-differentiated T-ALL cell lines. *Leukemia* **2014**; 28: 1145-8. doi: 10.1038/leu.2013.377.
18. Heo SK, Noh EK, Yu HM, Kim DK, Seo HJ, Lee YJ, et al. Radotinib enhances cytarabine (Ara-C)-induced acute myeloid leukemia cell death. *BMC Cancer* **2020**; 20: 1193. doi: 10.1186/s12885-020-07701-8.
19. Wu PS, Wang CY, Hsu HJ, Yen JH, Wu MJ. 8-Hydroxydaidzein induces apoptosis and inhibits AML-associated gene expression in U-937 cells: potential phytochemical for AML treatment. *Biomolecules* **2023**; 13: 1575. doi: 10.3390/biom13111575.
20. Zhou Y, Yu F, Luo B, Luo H, Liu C. Cytarabine (Ara-c) promotes cell apoptosis by inhibiting the phosphorylation of protein kinase B (AKT/PKB). *Process Biochem* **2019**; 82: 144-52. doi: 10.1016/j.procbio.2019.04.007.
21. Samareh Salavatipour M, Hoseinpour Kasgari F, Farsinejad A, Fatemi A, Mirzaee Khalilabadi R. Platelet-derived microparticles increase expression of hTERT in umbilical cord mesenchymal stem cells. *Res Mol Med* **2018**; 5: 31-40. doi: 10.18502/rmm.v5i4.3063.

22. Labrie A, Marshall A, Bedi H, Maurer-Spurej E. Characterization of platelet concentrates using dynamic light scattering. *Transfus Med Hemother* **2013**; 40: 93-100. doi: 10.1159/000350362.
23. Samareh Salavatipour M, Hoseinpour Kasgari F, Farsinejad A, Fatemi A, Hassanshahi G, Mirzaee Khalilabadi R. Platelet microparticles accelerate proliferation and growth of mesenchymal stem cells through longevity-related genes. *Arch Iran Med* **2021**; 24: 607-14. doi: 10.34172/aim.2021.86.
24. Aatonen MT, Ohman T, Nyman TA, Laitinen S, Grönholm M, Siljander PR. Isolation and characterization of platelet-derived extracellular vesicles. *J Extracell Vesicles* **2014**; 3: 24692. doi: 10.3402/jev.v3.24692.
25. Fatemi A, Kazemi A, Kashiri M, Safa M. Elevation of cAMP levels inhibits doxorubicin-induced apoptosis in pre-B ALL Nalm-6 cells through induction of BAD phosphorylation and inhibition of P53 accumulation. *Int J Mol Cell Med* **2015**; 4: 94-102.
26. Chou T, Martin N. PC Software and User's Guide: A Computer Program for Quantitation of Synergism and Antagonism in Drug Combinations, and the Determination of IC50, ED50 and LD50 Values. Paramus, NJ: ComboSyn; 2005.
27. Anani H, Baluchi I, Farsinejad A, Fatemi A, Mirzaee khalilabadi R. Betulinic acid exerts antitumor effects on acute promyelocytic leukemia cells possibly via hTERT downregulation. *Gene Rep* **2022**; 26: 101435. doi: 10.1016/j.genrep.2021.101435.
28. Anani H, Baluchi I, Farsinejad A, Fatemi A, Mirzaee Khalilabadi R. *Zataria multiflora* methanolic extract has antitumor properties on U266 multiple myeloma cell line. *Gene Rep* **2020**; 20: 100655. doi: 10.1016/j.genrep.2020.100655.
29. Lashkari M, Fatemi A, Mardani Valandani H, Mirzaee Khalilabadi R. Promising anti-leukemic effect of *Zataria multiflora* extract in combination with doxorubicin to combat acute lymphoblastic leukemia cells (Nalm-6) (in vitro and in silico). *Sci Rep* **2022**; 12: 12657. doi: 10.1038/s41598-022-16943-4.
30. Piktel D, Moore JC, Nesbit S, Sprowls SA, Craig MD, Rellick SL, et al. Chemotherapeutic activity of pitavastatin in vincristine resistant B-cell acute lymphoblastic leukemia. *Cancers (Basel)* **2023**; 15: 707. doi: 10.3390/cancers15030707.
31. Kanno S, Higurashi A, Watanabe Y, Shouji A, Asou K, Ishikawa M. Susceptibility to cytosine arabinoside (Ara-C)-induced cytotoxicity in human leukemia cell lines. *Toxicol Lett* **2004**; 152: 149-58. doi: 10.1016/j.toxlet.2004.04.014.
32. Chen Y, Gan D, Huang Q, Luo X, Lin D, Hu J. Emodin and its combination with cytarabine induce apoptosis in resistant acute myeloid leukemia cells in vitro and in vivo. *Cell Physiol Biochem* **2018**; 48: 2061-73. doi: 10.1159/000492544.
33. Buegry D, Wenz F, Groden C, Brockmann MA. Tumor-platelet interaction in solid tumors. *Int J Cancer* **2012**; 130: 2747-60. doi: 10.1002/ijc.27441.
34. D'Alessandro R, Messa C, Refolo MG, Carr BI. Modulation of sensitivity and resistance to multikinase inhibitors by microenvironmental platelet factors in HCC. *Expert Opin Pharmacother* **2015**; 16: 2773-80. doi: 10.1517/14656566.2015.1101065.
35. Huong PT, Nguyen LT, Nguyen XB, Lee SK, Bach DH. The role of platelets in the tumor-microenvironment and the drug resistance of cancer cells. *Cancers (Basel)* **2019**; 11: 240. doi: 10.3390/cancers11020240.
36. Pan Y, Wang Y, Wang Y, Xu S, Jiang F, Han Y, et al. Platelet-derived microvesicles (PMVs) in cancer progression and clinical applications. *Clin Transl Oncol* **2023**; 25: 873-81. doi: 10.1007/s12094-022-03014-3.
37. Pan Y, Liang H, Liu H, Li D, Chen X, Li L, et al. Platelet-secreted microRNA-223 promotes endothelial cell apoptosis induced by advanced glycation end products via targeting the insulin-like growth factor 1 receptor. *J Immunol* **2014**; 192: 437-46. doi: 10.4049/jimmunol.1301790.
38. Burnouf T, Chou ML, Lundy DJ, Chuang EY, Tseng CL, Goubran H. Expanding applications of allogeneic platelets, platelet lysates, and platelet extracellular vesicles in cell therapy, regenerative medicine, and targeted drug delivery. *J Biomed Sci* **2023**; 30: 79. doi: 10.1186/s12929-023-00972-w.
39. Indana HA, Puspitawati I, Mayasari DS, Hartopo AB. Association of acute hyperglycemia and diabetes mellitus with platelet-derived microparticle (PDMP) levels during acute myocardial infarction. *J ASEAN Fed Endocr Soc* **2023**; 38: 35-40. doi: 10.15605/jafes.038.02.03.
40. Ashoub MH, Samareh Salavatipour M, Hoseinpour Kasgari F, Mardani Valandani H, Mirzaee Khalilabadi R. Extracellular microvesicles: biologic properties, biogenesis, and applications in leukemia. *Mol Cell Biochem* **2024**; 479: 419-30. doi: 10.1007/s11010-023-04734-y.
41. Cacic D, Reikvam H, Nordgård O, Meyer P, Hervig T. Platelet microparticles protect acute myelogenous leukemia cells against daunorubicin-induced apoptosis. *Cancers (Basel)* **2021**; 13: 1870. doi: 10.3390/cancers13081870.
42. Gharib E, Veilleux V, Boudreau LH, Pichaud N, Robichaud GA. Platelet-derived microparticles provoke chronic lymphocytic leukemia malignancy through metabolic reprogramming. *Front Immunol* **2023**; 14: 1207631. doi: 10.3389/fimmu.2023.1207631.
43. Yafthian M, Yari F, Ghasemzadeh M, Fallah Azad V, Haghighi M. Induction of apoptosis in cancer cells of pre-B ALL patients after exposure to platelets, platelet-derived microparticles and soluble CD40 ligand. *Cell J* **2018**; 20: 120-6. doi: 10.22074/cellj.2018.5032.
44. Vismara M, Zarà M, Negri S, Canino J, Canobbio I, Barbieri SS, et al. Platelet-derived extracellular vesicles regulate cell cycle progression and cell migration in breast cancer cells. *Biochim Biophys Acta Mol Cell Res* **2021**; 1868: 118886. doi: 10.1016/j.bbamcr.2020.118886.
45. Michael JV, Wurtzel JGT, Mao GF, Rao AK, Kolpakov MA, Sabri A, et al. Platelet microparticles infiltrating solid tumors transfer miRNAs that suppress tumor growth. *Blood* **2017**; 130: 567-80. doi: 10.1182/blood-2016-11-751099.
46. Capriotti AL, Caruso G, Cavaliere C, Piovesana S, Samperi R, Laganà A. Proteomic characterization of human platelet-derived microparticles. *Anal Chim Acta* **2013**; 776: 57-63. doi: 10.1016/j.aca.2013.03.023.
47. Baj-Krzyworzeka M, Majka M, Pratico D, Ratajczak J, Vilaire G, Kijowski J, et al. Platelet-derived microparticles stimulate proliferation, survival, adhesion, and chemotaxis of hematopoietic cells. *Exp Hematol* **2002**; 30: 450-9. doi: 10.1016/s0301-472x(02)00791-9.
48. Cacic D, Reikvam H, Nordgård O, Meyer P, Hervig T. Platelet microparticles protect acute myelogenous leukemia cells against daunorubicin-induced apoptosis. *Cancers (Basel)* **2021**; 13: 1870. doi: 10.3390/cancers13081870.
49. Czabotar PE, Garcia-Saez AJ. Mechanisms of BCL-2 family proteins in mitochondrial apoptosis. *Nat Rev Mol Cell Biol* **2023**; 24: 732-48. doi: 10.1038/s41580-023-00629-4.
50. Velez J, Enciso LJ, Suarez M, Fiegl M, Grisimaldo A, López C, et al. Platelets promote mitochondrial uncoupling and resistance to apoptosis in leukemia cells: a novel paradigm for the bone marrow microenvironment. *Cancer Microenviron* **2014**; 7: 79-90. doi: 10.1007/s12307-014-0149-3.
51. Boilard E, Ducheux AC, Brisson A. The diversity of platelet microparticles. *Curr Opin Hematol* **2015**; 22: 437-44. doi: 10.1097/moh.0000000000000166.
52. Xiao BD, Zhao YJ, Jia XY, Wu J, Wang YG, Huang F. Multifaceted p21 in carcinogenesis, stemness of tumor and tumor therapy. *World J Stem Cells* **2020**; 12: 481-7. doi: 10.4252/wjsc.v12.i6.481.
53. Leão R, Apolónio JD, Lee D, Figueiredo A, Tabori U, Castelo-Branco P. Mechanisms of human telomerase reverse transcriptase (hTERT) regulation: clinical impacts in cancer. *J Biomed Sci* **2018**; 25: 22. doi: 10.1186/s12929-018-0422-8.
54. Nazari M, Javandoost E, Talebi M, Movassaghpour A, Soleimani M. Platelet microparticle controversial role in cancer. *Adv Pharm Bull* **2021**; 11: 39-55. doi: 10.34172/apb.2021.005.